arXiv:0904.0331v2 [math.QA] 23 Jan 2010

A MATRIX REALIZATION OF THE QUANTUM GROUP $\mathfrak{g}_{p,q}$

YUSUKE ARIKE

Abstract. In this paper we will find a matrix realizations of the quantum group $\mathfrak{g}_{p,q}$ defined in [4]. For this purpose, we construct all primitive idempotents and a basis of $\mathfrak{g}_{p_1,p_2}$. We determine the action of elements of the basis on the indecomposable projective modules, which give rise to a matrix realization of $\mathfrak{g}_{p_1,p_2}$. By using this result, we obtain a basis of the space of symmetric linear functions on $\mathfrak{g}_{p,q}$ and express the symmetric linear functions obtained by the left integral, the balancing element and the center of $\mathfrak{g}_{p,q}$ in term of this basis.

1. Introduction

In [2] and [3], it is pointed out that the triplet vertex operator algebra $\mathcal{W}(p)$ is closely related with the restricted quantum group $\overline{U}_q(sl_2)$ at the primitive $2p$-th root of the unity. It is also found in [2] and [3] find that the space of functions obtained by applying modular transformations to characters of simple modules of $\mathcal{W}(p)$ is isomorphic to the center of $\overline{U}_q(sl_2)$. It is conjectured in [3] that the category of modules of $\mathcal{W}(p)$ and the category of finite-dimensional modules of $\overline{U}_q(sl_2)$ are equivalent as tensor categories. It is proved the conjecture for $p = 2$. The equivalence of categories as abelian categories is proved in [10]. We also mention that the tensor category consisting of finite-dimensional $\overline{U}_q(sl_2)$-modules is not braided if $p \neq 2$ ([7]).

Feigin, Gainutdinov, Semikhatov and Tipunin also construct the vertex operator algebra $\mathcal{W}(p_1, p_2)$ for coprime positive integers p_1 and p_2. They find $2p_1p_2 + \frac{1}{2}(p_1 - 1)(p_2 - 1)$ non-isomorphic simple modules and show that the space of functions obtained by applying modular transformations on characters of simple modules is $\frac{1}{2}(3p_1 - 1)(3p_2 - 1)$-dimensional. They study in [4] the unimodular finite-dimensional Hopf algebra $\mathfrak{g}_{p_1,p_2}$ which is a quotient algebra of the tensor product of two restricted quantum groups $\overline{U}_{q_1}(sl_2)$ at $q_1 = \exp(\sqrt{-1}p_2\pi/p_1)$ and $\overline{U}_{q_2}(sl_2)$ at $q_1 = \exp(\sqrt{-1}p_1\pi/p_2)$. In [5], they show that $\mathfrak{g}_{p_1,p_2}$ has $2p_1p_2$ non-isomorphic simple modules, which shows that the category of modules of $\mathcal{W}(p_1, p_2)$ is not equivalent to the category of finite-dimensional modules of $\mathfrak{g}_{p_1,p_2}$. They prove that the space of functions obtained by applying modular transformations to characters of simple modules of $\mathcal{W}(p_1, p_2)$ and the center of $\mathfrak{g}_{p_1,p_2}$, on which the group $SL_2(\mathbb{Z})$ naturally acts, are equivalent as representations of $SL_2(\mathbb{Z})$.

In this paper, we determine a matrix realization of $\mathfrak{g}_{p_1,p_2}$. More precisely, it is shown in [5] that the quantum group $\mathfrak{g}_{p_1,p_2}$ has a decomposition into two-sided ideals

$$\mathfrak{g}_{p_1,p_2} = Q(p_1, p_2) \oplus Q(0, p_2) \oplus \bigoplus_{r_1=1}^{p_1-1} Q(r_1, p_2) \oplus \bigoplus_{r_2=1}^{p_2-1} Q(p_1, r_2) \oplus \bigoplus_{(r_1,r_2)\in I} Q(r_1, r_2), \tag{1.0.1}$$

2000 *Mathematics Subject Classification.* Primary 16W35, Secondary 17B37, 81R05.

where $I = \{(r_1, r_2) | 1 \leq r_i \leq p_i - 1,\ p_1r_2 + p_2r_1 \leq p_1p_2\}$, we will get a matrix realization of a two-sided ideal which appears in (1.0.1). In order to construct a basis, we first determine primitive idempotents and derive direct sum decompositions of two-sided ideals into their indecomposable left ideals as it is done in [1] and [12]. Then, by calculating actions of two-sided ideals on projective modules, we can have matrix realizations of two-sided ideals.

Theorem 6.1. *The algebras $Q(p_1, p_2)$ and $Q(0, p_2)$ are isomorphic to the matrix algebras $M_{p_1p_2}(\mathbb{C})$.*

Theorem 6.2. *For $1 \leq r_1 \leq p_1 - 1$, the algebra $Q(r_1, p_2)$ is isomorphic to the subalgebra of $M_{2p_1p_2}(\mathbb{C}) \oplus M_{2p_1p_2}(\mathbb{C})$ with the shape:*

$$\left(\begin{bmatrix} B^{+,\uparrow}_{r_1,p_2} & 0 & 0 & 0 \\ B^{+,\leftarrow}_{r_1,p_2} & B^{-,\uparrow}_{p_1-r_1,p_2} & 0 & 0 \\ B^{+,\rightarrow}_{r_1,p_2} & 0 & B^{-,\uparrow}_{p_1-r_1,p_2} & 0 \\ B^{+,\downarrow}_{r_1,p_2} & B^{-,\rightarrow}_{p_1-r_1,p_2} & B^{-,\leftarrow}_{p_1-r_1,p_2} & B^{+,\uparrow}_{r_1,p_2} \end{bmatrix}, \begin{bmatrix} B^{-,\uparrow}_{p_1-r_1,p_2} & 0 & 0 & 0 \\ B^{-,\leftarrow}_{p_1-r_1,p_2} & B^{+,\uparrow}_{r_1,p_2} & 0 & 0 \\ B^{-,\rightarrow}_{p_1-r_1,p_2} & 0 & B^{+,\uparrow}_{r_1,p_2} & 0 \\ B^{-,\downarrow}_{p_1-r_1,p_2} & B^{+,\rightarrow}_{r_1,p_2} & B^{-,\leftarrow}_{r_1,p_2} & B^{-,\uparrow}_{p_1-r_1,p_2} \end{bmatrix}\right).$$

Theorem 6.3. *For $1 \leq r_1 \leq p_1 - 1$, the algebra $Q(r_1, p_2)$ is isomorphic to the subalgebra of $M_{2p_1p_2}(\mathbb{C}) \oplus M_{2p_1p_2}(\mathbb{C})$:*

$$\left(\begin{bmatrix} B^{+,\uparrow}_{p_1,r_2} & 0 & 0 & 0 \\ B^{+,\leftarrow}_{p_1,r_2} & B^{-,\uparrow}_{p_1,p_2-r_2} & 0 & 0 \\ B^{+,\rightarrow}_{p_1,r_2} & 0 & B^{-,\uparrow}_{p_1,p_2-r_2} & 0 \\ B^{+,\downarrow}_{p_1,r_2} & B^{-,\rightarrow}_{p_1,p_2-r_2} & B^{-,\leftarrow}_{p_1,p_2-r_2} & B^{+,\uparrow}_{p_1,r_2} \end{bmatrix}, \begin{bmatrix} B^{-,\uparrow}_{p_1,p_2-r_2} & 0 & 0 & 0 \\ B^{-,\leftarrow}_{p_1,p_2-r_2} & B^{+,\uparrow}_{p_1,r_2} & 0 & 0 \\ B^{-,\rightarrow}_{p_1,p_2-r_2} & 0 & B^{+,\uparrow}_{p_1,r_2} & 0 \\ B^{-}_{p_1,p_2-r_2} & B^{+,\rightarrow}_{p_1,r_2} & B^{+,\leftarrow}_{p_1,r_2} & B^{-,\uparrow}_{p_1,p_2-r_2} \end{bmatrix}\right).$$

Theorem 6.4. *The algebra of $Q(r_1, r_2)$ with $(r_1, r_2) \in I$ is isomorphic to the subalgebra of $M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C})$ given by*

$$\left(\begin{bmatrix} T^{+}_{r_1,r_2} & 0 & 0 & 0 \\ L^{+}_{r_1,r_2} & T^{-}_{r_1,p_2-r_2} & 0 & 0 \\ R^{+}_{r_1,r_2} & 0 & T^{-}_{r_1,p_2-r_2} & 0 \\ B^{+}_{r_1,r_2} & R^{-}_{r_1,p_2-r_2} & L^{-}_{r_1,p_2-r_2} & T^{+}_{r_1,r_2} \end{bmatrix}, \begin{bmatrix} T^{-}_{p_1-r_1,r_2} & 0 & 0 & 0 \\ L^{-}_{p_1-r_1,r_2} & T^{+}_{p_1-r_1,p_2-r_2} & 0 & 0 \\ R^{-}_{p_1-r_1,r_2} & 0 & T^{+}_{p_1-r_1,p_2-r_2} & 0 \\ B^{-}_{p_1-r_1,r_2} & R^{+}_{p_1-r_1,p_2-r_2} & L^{+}_{p_1-r_1,p_2-r_2} & T^{-}_{p_1-r_1,r_2} \end{bmatrix},\right.$$

$$\left.\begin{bmatrix} T^{-}_{r_1,p_2-r_2} & 0 & 0 & 0 \\ L^{-}_{r_1,p_2-r_2} & T^{+}_{r_1,r_2} & 0 & 0 \\ R^{-}_{r_1,p_2-r_2} & 0 & T^{+}_{r_1,r_2} & 0 \\ B^{-}_{r_1,p_2-r_2} & R^{+}_{r_1,r_2} & L^{+}_{r_1,r_2} & T_{r_1,p_2-r_2} \end{bmatrix}, \begin{bmatrix} T^{+}_{p_1-r_1,p_2-r_2} & 0 & 0 & 0 \\ L^{+}_{p_1-r_1,p_2-r_2} & T^{-}_{p_1-r_1,r_2} & 0 & 0 \\ R^{+}_{p_1-r_1,p_2-r_2} & 0 & T^{-}_{p_1-r_1,r_2} & 0 \\ B^{+}_{p_1-r_1,p_2-r_2} & R^{-}_{p_1-r_1,r_2} & L^{-}_{p_1-r_1,r_2} & T^{+}_{p_1-r_1,p_2-r_2} \end{bmatrix}\right)$$

where

$$X^{\alpha}_{r_1,r_2} = \begin{bmatrix} X^{\alpha,\uparrow}_{r_1,r_2} & 0 & 0 & 0 \\ X^{\alpha,\leftarrow}_{r_1,r_2} & X^{-\alpha,\uparrow}_{p_1-r_1,r_2} & 0 & 0 \\ X^{\alpha,\rightarrow}_{r_1,r_2} & 0 & X^{-\alpha,\uparrow}_{p_1-r_1,r_2} & 0 \\ X^{\alpha,\downarrow}_{r_1,r_2} & X^{-\alpha,\rightarrow}_{p_1-r_1,r_2} & X^{-\alpha,\leftarrow}_{p_1-r_1,r_2} & X^{\alpha,\uparrow}_{r_1,r_2} \end{bmatrix}$$

and $X = T, R, L, B$.

By using these matrix realizations, we will get a basis of the space of symmetric linear functions $\mathrm{SLF}(\boldsymbol{\mathfrak{g}_{p_1,p_2}})$ as the sums of traces of matrix blocks appeared in the matrix realizations.

As it is shown in [11], The space $\mathrm{SLF}(\mathfrak{g}_{p_1,p_2})$ is isomorphic to the center of $\mathfrak{g}_{p_1,p_2}$. This isomorphism is given by $c \mapsto t^{-1}c \rightharpoonup \lambda$ where λ is the left integral and t is an invertible element in $\mathfrak{g}_{p_1,p_2}$ such that $S^2(x) = txt^{-1}$ for all $x \in \mathfrak{g}_{p_1,p_2}$. Then we can determine the relations between the basis of symmetric linear functions and symmetric linear functions determined by the action of the central elements and balancing element of $\mathfrak{g}_{p,q}$ on integrals.

In [5], the space of q-characters of $\mathfrak{g}_{p_1,p_2}$ is determined. The most important thing is that they construct a basis of the space of q-characters. Since the space of the q-characters is $\{\beta \in {\mathfrak{g}_{p_1,p_2}}^* | \beta(xy) = \beta(S^2(y)x),\ x, y \in \mathfrak{g}_{p_1,p_2}\}$ (c.f. [5]) and the square of the antipode of $\mathfrak{g}_{p_1,p_2}$ is an inner automorphism by the balancing element g of $\mathfrak{g}_{p_1,p_2}$ (see [5]), any symmetric linear function is given by $\beta \leftharpoonup g$ where β is a q-character. Under this correspondence, any element of our basis of $\mathrm{SLF}(\mathfrak{g}_{p_1,p_2})$ is mapped to a scalar multiple of an element of the basis of the space of q-characters constructed in [5] (see Appendix A).

This paper is organized as follows. In section 2, we recall the basic definitions and properties of symmetric linear functions and integrals of Hopf algebras. In section 3, we recall the definition of the algebra $\mathfrak{g}_{p_1,p_2}$ and its integrals and balancing element given in [5]. We also introduce the list of non-isomorphic simple modules $\mathfrak{g}_{p_1,p_2}$ given in [5]. In section 4, we construct indecomposable modules as left ideals of $\mathfrak{g}_{p_1,p_2}$. In section 5, we find the primitive idempotents of $\mathfrak{g}_{p_1,p_2}$ and show that the indecomposable modules constructed in section 4 give rise to the list of non-isomorphic indecomposable projective modules. In addition, we give a decomposition of $\mathfrak{g}_{p_1,p_2}$ into subalgebras. In section 6, we determine the matrix realization of each subalgebra of $\mathfrak{g}_{p_1,p_2}$ as the subalgebras of direct sum of matrix algebras. In section 7, we construct the symmetric linear functions on $\mathfrak{g}_{p_1,p_2}$ which form a basis of the space of symmetric linear functions on $\mathfrak{g}_{p_1,p_2}$. We also determine the relations between the basis and the symmetric linear functions given by the left integral, the balancing element and the central elements. In Appendix, we give a correspondence between the basis of q-characters of $\mathfrak{g}_{p_1,p_2}$ obtained in [5] and the basis of symmetric linear functions on $\mathfrak{g}_{p_1,p_2}$

Acknowledgment

The author is grateful to K. Nagatomo for his continuous encouragement and useful comments.

2. Preliminaries

In this paper we will always work over the complex number field $\mathbb{C}$. For any vector space V we denote its dual space $\mathrm{Hom}_{\mathbb{C}}(V, \mathbb{C})$ by V^*.

2.1. **Symmetric linear functions.**

Let A be a finite-dimensional associative algebra. A symmetric linear function φ on A is an element of A^* which satisfies $\varphi(ab) = \varphi(ba)$ for all $a, b \in A$. We denote the space of symmetric linear functions on A by $\mathrm{SLF}(A)$. If A is a finite-dimensional Hopf algebra, the space $\mathrm{SLF}(A)$ coincides with the space of cocommutative elements of A^* (see [11]).

2.2. **Integrals and the square of antipode of Hopf algebras.** Let A be a finite-dimensional Hopf algebra with the coproduct Δ, the counit ε and the antipode S. Any element of the subspaces

$$\mathcal{L}_A = \{\Lambda \in A \mid a\Lambda = \varepsilon(a)\Lambda \text{ for all } a \in A\},$$
$$\mathcal{R}_A = \{\Lambda \in A \mid \Lambda a = \varepsilon(a)\Lambda \text{ for all } a \in A\},$$

is called a left integral and a right integral of A, respectively. Since A is finite-dimensional, the space $\mathcal{L}_A$ (respectively $\mathcal{R}_A$) is one-dimensional (cf. [9]). Similarly a left (respectively, right) integral of the dual Hopf algebra A^* is an element $\lambda \in A^*$ which satisfies $p\lambda = p(1)\lambda$ (respectively, $\lambda p = p(1)\lambda$) for all $p \in A^*$. Equivalently we can see

$$\mathcal{L}_{A^*} = \{\lambda \in A^* \mid (1 \otimes \lambda)\Delta(x) = \lambda(x) \text{ for all } x \in A\},$$
$$\mathcal{R}_{A^*} = \{\lambda \in A^* \mid (\lambda \otimes 1)\Delta(x) = \lambda(x) \text{ for all } x \in A\}.$$

If $\mathcal{L}_A = \mathcal{R}_A$ the Hopf algebra A is called *unimodular*.

Proposition 2.1 ([11]). *Let A be a finite-dimensional unimodular Hopf algebra with the antipode S. Suppose that λ is a left integral of A^* and that μ is a right integral of A^*. Then*

(1) $\lambda(ab) = \lambda(bS^2(a))$,
(2) $\mu(ab) = \mu(S^2(b)a)$.

The square of the antipode is called *inner* if there exists an invertible element t such that $S^2(x) = txt^{-1}$ for all $x \in A$.

Denote by $\rightharpoonup$ and $\leftharpoonup$ the left and right actions of A on A^*:

$$a \rightharpoonup p(b) = p(ba),\ p \leftharpoonup a(b) = p(ab),$$

for $p \in A^*$ and $a, b \in A$.

Proposition 2.2 ([11]). *Let A be a finite-dimensional unimodular Hopf algebra with the antipode S. Suppose that λ is a left integral of A^* and that μ is a right integral of A^*. If S^2 is inner, the linear maps f_ℓ and $f_r : Z(A) \to \mathrm{SLF}(A)$ defined by $f_\ell(c) = t^{-1}c \rightharpoonup \lambda$ and $f_r(c) = \mu \leftharpoonup tc$ are isomorphisms where $Z(A)$ denote the center of A.*

3. THE HOPF ALGEBRA $\mathfrak{g}_{p,q}$

3.1. **Definition.** Let p_1 and p_2 be coprime positive integers. Set $q = \exp(\frac{\pi\sqrt{-1}}{2p_1p_2})$, $q_1 = q^{2p_2} = \exp(\frac{\pi\sqrt{-1}}{p_1})$, and $q_2 = q^{2p_1} = \exp(\frac{\pi\sqrt{-1}}{p_2})$. We use the q-integers and q-binomial coefficients

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}},\ \begin{bmatrix} m \\ n \end{bmatrix}_q = \frac{[m]_q!}{[n]_q![m-n]_q!},\ [n]_q! = [n]_q \cdots [2]_q[1]_q.$$

We also define

$$[m]_1 = [m]_{q_1^{p_2}},\ [m]_2 = [m]_{q_2^{p_1}},\ \begin{bmatrix} m \\ n \end{bmatrix}_1 = \begin{bmatrix} m \\ n \end{bmatrix}_{q_1^{p_2}},\ \begin{bmatrix} m \\ n \end{bmatrix}_2 = \begin{bmatrix} m \\ n \end{bmatrix}_{q_2^{p_1}}.$$

Note that

$$[p_1 - m]_1 = (-1)^{p_2+1}[m]_1 \text{ and } [p_2 - m]_2 = (-1)^{p_1+1}[m]_2.$$

$\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ is a Hopf algebra over $\mathbb{C}$ generated by e_i, f_i and $K^{\pm}$ for $i = 1, 2$ with relations

$$\begin{aligned}
&KK^{-1} = K^{-1}K = 1,\\
&e_i^{p_i} = f_i^{p_i} = 0,\ K^{2p_1p_2} = 1,\ (i = 1, 2),\\
&Ke_iK^{-1} = q_i^2 e_i,\ Kf_iK^{-1} = q_i^{-2} f_i,\ (i = 1, 2),\\
&e_1e_2 = e_2e_1,\ f_1f_2 = f_2f_1,\\
&[e_1, f_1] = \frac{K^{p_2} - K^{-p_2}}{q_1^{p_2} - q_1^{-p_2}},\ [e_2, f_2] = \frac{K^{p_1} - K^{-p_1}}{q_2^{p_1} - q_2^{-p_1}},
\end{aligned}$$

as an algebra. The coproduct Δ, counit ε, and antipode S are given by

$$\begin{aligned}
&\Delta(e_1) = e_1 \otimes 1 + K^{p_2} \otimes e_1,\ \Delta(e_2) = e_2 \otimes K^{p_1} + 1 \otimes e_2,\\
&\Delta(f_1) = f_1 \otimes K^{-p_2} + 1 \otimes f_1,\ \Delta(f_2) = f_2 \otimes 1 + K^{-p_1} \otimes f_2,\\
&\Delta(K) = K \otimes K,\ \varepsilon(e_i) = \varepsilon(f_i) = 0,\ \varepsilon(K) = K,\\
&S(e_1) = -K^{-p_2}e_1,\ S(e_2) = -e_2K^{-p_1},\\
&S(f_1) = -f_1K^{p_2},\ S(f_2) = -K^{p_1}f_2,\ S(K) = K^{-1}.
\end{aligned}$$

Proposition 3.1. *The* $2p_1^3p_2^3$ *elements* $e_1^{m_1}e_2^{m_2}f_1^{n_1}f_2^{n_2}K^{\ell}$*, where* $0 \le m_i, n_i \le p_i - 1$ *and* $0 \le \ell \le 2p_1p_2 - 1$*, form a basis of* $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ *as a vector space over* $\mathbb{C}$*.*

In [5], it is shown that the restricted quantum groups $\overline{U}_{q_1^{p_2}}(sl_2) = \langle e_1, f_1, K_1\rangle$ and $\overline{U}_{q_2^{p_1}}(sl_2) = \langle e_2, f_2, K_2\rangle$ are embedded in $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ by

$$\begin{aligned}
&e_1 \mapsto e_1,\ f_1 \mapsto f_1,\ K_1 \mapsto K^{p_2},\\
&e_2 \mapsto e_2,\ f_2 \mapsto f_2,\ K_2 \mapsto K^{p_1}
\end{aligned}$$

and that

$$\boldsymbol{\mathfrak{g}_{p_1,p_2}} \cong \overline{U}_{q_1^{p_2}}(sl_2) \otimes \overline{U}_{q_2^{p_1}}(sl_2)/(K_1^{p_2} \otimes 1 - 1 \otimes K_2^{p_2}),$$

where $(K_1^{p_1} \otimes 1 - 1 \otimes K_2^{p_2})$ is the Hopf ideal generated by $(K_1^{p_1} \otimes 1 - 1 \otimes K_2^{p_2})$.

3.2. **Integrals and the square of the antipode.**

The space of left and right integrals in $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ and the space of right integrals on $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ are determined in [5]. The space spanned by

$$e_1^{p_1-1}e_2^{p_2-1}f_1^{p_1-1}f_2^{p_2-1}\sum_{\ell=0}^{2p_1p_2-1} K^{\ell},$$

coincides with the space of left integrals of $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$, which also coincides with the space of right integrals of $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$. This shows that $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ is unimodular.

Define the linear functions on $\boldsymbol{\mathfrak{g}_{p_1,p_2}}$ by

$$\lambda(e_1^{m_1}e_2^{m_2}f_1^{n_1}f_2^{n_2}K^{\ell}) = \delta_{m_1,p_1-1}\delta_{m_2,p_2-1}\delta_{n_1,p_1-1}\delta_{n_2,p_2-1}\delta_{\ell,p_2-p_1}, \tag{3.2.1}$$

$$\mu(e_1^{m_1}e_2^{m_2}f_1^{n_1}f_2^{n_2}K^{\ell}) = \delta_{m_1,p_1-1}\delta_{m_2,p_2-1}\delta_{n_1,p_1-1}\delta_{n_2,p_2-1}\delta_{\ell,p_1-p_2}, \tag{3.2.2}$$

for $0 \le m_i \le p_i - 1$ and $-(p_1p_2 - 1) \le \ell \le p_1p_2$. Using the induction, we can see that

$$\begin{aligned}\Delta(e_1^{m_1}e_2^{m_2}f_1^{n_1}f_2^{n_2}K^{\ell}) &= \sum_{r_1=0}^{m_1}\sum_{r_2=0}^{m_2}\sum_{s_1=0}^{n_1}\sum_{s_2=0}^{n_2} q_1^{p_2(m_1-r_1)+p_2s_1(n_1-r_1)-2p_2s_1(m_1-r_1)} \\ &\times q_2^{p_1r_2(m_2-r_2)+p_1s_2(n_2-s_2)-2p_1r_2(n_2-s_2)} \begin{bmatrix} m_1 \\ r_1 \end{bmatrix}_1 \begin{bmatrix} m_2 \\ r_2 \end{bmatrix}_2 \begin{bmatrix} n_1 \\ s_1 \end{bmatrix}_1 \begin{bmatrix} n_2 \\ s_2 \end{bmatrix}_2 \\ &\times e_1^{r_1}e_2^{r_2}f_1^{s_1}f_2^{s_2}K^{p_2(m_1-r_1)-p_1(n_2-s_2)+\ell} \otimes e_1^{m_1-r_1}e_2^{m_2-r_2}f_1^{n_1-s_1}f_2^{n_2-s_2}K^{p_1r_2-p_2s_1+\ell}.\end{aligned}$$

Proposition 3.2 ([5]). *Each of the spaces of left and right integrals in* $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ *is spanned by* λ *and* μ, *respectively.*

It is also shown in [5] that the square of the antipode of $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ is inner and that the balancing element of $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ is given as follows.

Proposition 3.3 ([5]). *For any* $x \in \mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$, *we have* $S^2(x) = gxg^{-1}$ *where* $g = K^{p_1-p_2}$.

3.3. **Simple modules.** The algebra $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ has $2p_1p_2$ non-isomorphic simple modules (see [5]). Let $\{X^{\alpha}_{r_1,r_2} \mid \alpha = \pm, 1 \le r_1 \le p_1, 1 \le r_2 \le p_2\}$ be the complete list of non-isomorphic simple modules of $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$. The simple module $X^{\alpha}_{r_1,r_2}$ has a basis formed by weight vectors $\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2)$, $0 \le n_1 \le r_1 - 1$ and $0 \le n_2 \le r_2 - 1$ with the action of $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ defined by

$$K\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2) = \alpha q_1^{r_1-1-2n_1}q_2^{r_2-1-2n_2}\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2),$$

$$e_1\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2) = \begin{cases} \varphi_1^{\alpha}(n_1,r_1,r_2)\mathsf{b}^{\alpha}_{n_1-1,n_2}(r_1,r_2), & n_1 \ne 0, \\ 0, & n_1 = 0, \end{cases}$$

$$e_2\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2) = \begin{cases} \varphi_2^{\alpha}(n_2,r_1,r_2)\mathsf{b}^{\alpha}_{n_1,n_2-1}(r_1,r_2), & n_2 \ne 0, \\ 0, & n_2 = 0, \end{cases}$$

$$f_1\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2) = \begin{cases} \mathsf{b}^{\alpha}_{n_1+1,n_2}(r_1,r_2), & n_1 \ne r_1 - 1, \\ 0, & n_1 = r_1 - 1, \end{cases}$$

$$f_2\mathsf{b}^{\alpha}_{n_1,n_2}(r_1,r_2) = \begin{cases} \mathsf{b}^{\alpha}_{n_1,n_2+2}(r_1,r_2), & n_2 \ne r_2 - 1, \\ 0, & n_2 = r_2 - 1, \end{cases}$$

where

$$\varphi_1^{\alpha}(n_1,r_1,r_2) = \alpha^{p_2}(-1)^{r_2-1}[n_1]_1[r_1-n_1]_1, \tag{3.3.1}$$

$$\varphi_2^{\alpha}(n_2,r_1,r_2) = \alpha^{p_1}(-1)^{r_1-1}[n_2]_2[r_2-n_2]_2, \tag{3.3.2}$$

for $1 \le n_i \le r_i - 1$. We note that

$$\varphi_1^{-\alpha}(k_1,p_1-r_1,r_2) = \varphi_1^{\alpha}(k_1,p_1-r_1,p_2-r_2), \tag{3.3.3}$$

$$\varphi_1^{\alpha}(n_1,r_1,r_2) = \varphi_1^{-\alpha}(n_1,r_1,p_2-r_2), \tag{3.3.4}$$

for $1 \le n_1 \le r_1 - 1$ and $1 \le k_1 \le p_1 - r_1 - 1$, and that

$$\varphi_2^{-\alpha}(k_2,r_1,p_2-r_2) = \varphi_2^{\alpha}(k_2,p_1-r_1,p_2-r_2), \tag{3.3.5}$$

$$\varphi_2^{\alpha}(n_2,r_1,r_2) = \varphi_2^{-\alpha}(n_1,p_1-r_1,r_2), \tag{3.3.6}$$

for $1 \le n_2 \le r_2 - 1$ and $1 \le k_2 \le p_2 - r_2 - 1$.

4. CONSTRUCTION OF INDECOMPOSABLE LEFT IDEALS

In this section we construct left ideals which are isomorphic to $P^{\alpha}_{r_1,r_2}$ whose socle is isomorphic to a simple module $X^{\alpha}_{r_1,r_2}$ for $(r_1, r_2) \neq (p_1, p_2)$. The structure of module $P^{\alpha}_{r_1,r_2}$ is described in [5].

4.1. **Highest weight vectors in $\mathfrak{g}_{p_1,p_2}$.** For $1 \leq r_i \leq p_i$ and $1 \leq s_i \leq r_i$, let us set

$$v^{\alpha}_{r_1,r_2}(s_1, s_2) = \sum_{\ell=0}^{2p_1p_2-1} (\alpha q_1^{-(r_1-2s_1+1)} q_2^{-(r_2-2s_2+1)})^{\ell} K^{\ell},$$

and

$$b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) = f_1^{n_1} f_2^{n_2} e_1^{p_1-1} e_2^{p_2-1} f_1^{p_1-s_1} f_2^{p_2-s_2} v^{\alpha}_{r_1,r_2}(s_1, s_2),$$

for $0 \leq n_i \leq r_i - 1$. We then see that

$$K b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) = \alpha q_1^{r_1-1-2n_1} q_2^{r_2-1-2n_2} b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) \tag{4.1.1}$$

for $0 \leq n_1 \leq r_1 - 1$ and $0 \leq n_2 \leq r_2 - 1$. We also have

$$f_1 b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) = b^{\alpha,\downarrow}_{n_1+1,n_2}(r_1, r_2, s_1, s_2), \tag{4.1.2}$$

$$f_2 b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) = b^{\alpha,\downarrow}_{n_1,n_2+1}(r_1, r_2, s_1, s_2) \tag{4.1.3}$$

for $0 \leq n_1 \leq r_1 - 2$ and $0 \leq n_2 \leq r_2 - 2$.

Lemma 4.1 ([6])**.** *For $1 \leq n_i \leq p_i - 1$, $i, j \in \{1, 2\}$ and $i \neq j$, the following relations hold in $\mathfrak{g}_{p_1,p_2}$:*

$$\begin{aligned}
[e_i, f_i^{n_i}] &= [n_i]_i f_i^{n_i-1} \frac{q_i^{-p_j(n_i-1)} K^{p_j} - q_i^{p_j(n_i-1)} K^{-p_j}}{q_i^{p_j} - q_i^{-p_j}} \\
&= [n_i]_i \frac{q_i^{p_j(n_i-1)} K^{p_j} - q_i^{-p_j(n_i-1)} K^{-p_j}}{q_i^{p_j} - q_i^{-p_j}} f_i^{n_i-1}, \\
[e_i^{n_i}, f_i] &= [n_i]_i e_i^{n_i-1} \frac{q_i^{p_j(n_i-1)} K^{p_j} - q_i^{-p_j(n_i-1)} K^{-p_j}}{q_i^{p_j} - q_i^{-p_j}} \\
&= [n_i]_i \frac{q_i^{-p_j(n_i-1)} K^{p_j} - q_i^{p_j(n_i-1)} K^{-p_j}}{q_i^{p_j} - q_i^{-p_j}} e_i^{n_i-1}.
\end{aligned}$$

Lemma 4.2. *We have*

$$\begin{aligned}
&e_1 b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} \varphi_1^{\alpha}(n_1, r_1, r_2) b^{\alpha,\downarrow}_{n_1-1,n_2}(r_1, r_2, s_1, s_2), & 1 \leq n_1 \leq r_1 - 1, \\ 0, & n_1 = 0, \end{cases}
\end{aligned} \tag{4.1.4}$$

$$\begin{aligned}
&e_2 b^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} \varphi_2^{\alpha}(n_2, r_1, r_2) b^{\alpha,\downarrow}_{n_1,n_2-1}(r_1, r_2, s_1, s_2), & 1 \leq n_2 \leq r_2 - 1, \\ 0, & n_2 = 0, \end{cases}
\end{aligned} \tag{4.1.5}$$

Proof. By Lemma 4.1, we obtain

$$\begin{aligned}
&e_1 b^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2)\\
=&e_1 f_1^{n_1} f_2^{n_2} e_1^{p_1-1} e_2^{p_2-1} f_1^{p_1-s_1} f_2^{p_2-s_2} v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
=&[n_1]_1 \frac{q_1^{p_2(n_1-1)}K^{p_2} - q_1^{-p_2(n_1-1)}K^{-p_2}}{q_1^{p_2}-q_1^{-p_2}} b^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2)\\
=&\alpha^{p_2}(-1)^{r_2-1}[n_1]_1[r_1-n_1] b^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2),
\end{aligned}$$

for $1 \le n_1 \le r_1 - 1$. Similarly we have (4.1.5). □

The space spanned by $b^{\alpha}_{n_1,n_2}(r_1,r_2,s_1,s_2)$ with $0 \le n_1 \le r_1-1$ and $0 \le n_2 \le r_2 - 1$ is isomorphic to the simple module $X^{\alpha}_{r_1,r_2}$. This fact will be proved in next subsection.

Proposition 4.3. (1) *For* $1 \le r_1 \le p_1 - 1$,

$$\begin{aligned}
&b^{\alpha,\downarrow}_{0,n_2}(r_1,r_2,s_1,s_2)\\
=&f_1 f_2^{n_2} \sum_{m_1=1}^{p_1-r_1} \gamma^{\alpha}_{m_1}(r_1,r_2) e_1^{p_1-m_1} e_2^{p_2-1} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2} v^{\alpha}_{r_1,r_2}(s_1,s_2),
\end{aligned}$$

where

$$\gamma^{\alpha}_{m_1}(r_1,r_2) = \prod_{k_1=p_1-r_1-(m_1-1)}^{p_1-r_1-1} \varphi_1^{-\alpha}(k_1, p_1-r_1, r_2).$$

(2) *For* $1 \le r_2 \le p_2 - 1$,

$$\begin{aligned}
&b^{\alpha,\downarrow}_{n_1,0}(r_1,r_2,s_1,s_2)\\
=&f_1^{n_1} f_2 \sum_{m_2=1}^{p_2-r_2} \delta^{\alpha}_{m_2}(r_1,r_2) e_1^{p_1-1} e_2^{p_2-m_2} f_1^{p_1-s_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{r_1,r_2}(s_1,s_2),
\end{aligned}$$

where

$$\delta^{\alpha}_{m_2}(r_1,r_2) = \prod_{k_2=p_2-r_2-(m_2-1)}^{p_2-r_2-1} \varphi_2^{-\alpha}(k_2, r_1, p_2-r_2).$$

(3) *For* $1 \le r_1 \le p_1 - 1$ *and* $1 \le r_2 \le p_2 - 1$,

$$\begin{aligned}
b^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2) =& f_1 f_2 \sum_{m_1=1}^{p_1-r_1} \sum_{m_2=1}^{p_2-r_2} \gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2)\\
&\times e_1^{p_1-m_1} e_2^{p_2-m_2} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{r_1,r_2}(s_1,s_2).
\end{aligned}$$

Proof. We only prove (1). proofs of others are similar.

By Lemma 4.1, we have

$$\begin{aligned}
&f_1e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
=&e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1+1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
&\qquad-[e_1^{p_1-m_1},f_1]e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
=&e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1+1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
&-\varphi_1^{-\alpha}(m_1,p_1-r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2).
\end{aligned}$$

Thus we have

$$\begin{aligned}
&f_1\sum_{m_1=1}^{p_1-r_1}\gamma^{\alpha}_{m_1}(r_1,r_2)e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
=&\sum_{m_1=1}^{p_1-r_1}\gamma^{\alpha}_{m_1}(r_1,r_2)e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1+1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
&-\sum_{m_1=1}^{p_1-r_1-1}\gamma^{\alpha}_{m_1+1}(r_1,r_2)e_1^{p_1-m_1-1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2)\\
&=e_1^{p_1-1}e_2^{p_2-1}f_1^{p_1-s_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(s_1,s_2).
\end{aligned}$$

□

Set

$$\begin{aligned}
\Phi^{\alpha}(r_1,r_2)=2p_1p_2&\prod_{i_1=1}^{r_1-1}\varphi_1^{\alpha}(i_1,r_1,r_2)\prod_{i_2=1}^{r_2-1}\varphi_2^{\alpha}(k_2,r_1,r_2)\\
&\times\prod_{k_1=1}^{p_1-r_1-1}\varphi_1^{-\alpha}(k_1,p_1-r_1,r_2)\prod_{k_2=1}^{p_2-r_2-1}\varphi_2^{-\alpha}(k_2,r_1,p_2-r_2),\\
\Psi_1^{\alpha}(r_1,r_2)=&\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_2)}+\sum_{j_1=1}^{p_1-r_1-1}\frac{1}{\varphi_1^{-\alpha}(j_1,p_1-r_1,r_2)},\\
\Psi_2^{\alpha}(r_1,r_2)=&\sum_{i_2=1}^{r_2-1}\frac{1}{\varphi_2^{\alpha}(i_2,r_1,r_2)}+\sum_{j_2=1}^{p_2-r_2-1}\frac{1}{\varphi_2^{-\alpha}(j_2,r_1,p_2-r_2)}.
\end{aligned}$$

Then we can see that $\Phi^{\alpha}(r_1,r_2)=\Phi^{-\alpha}(p_1-r_1,r_2)=\Phi^{-\alpha}(r_1,p_2-r_2)=\Phi^{\alpha}(p_1-r_1,p_2-r_2)$ and that $\Psi_i^{\alpha}(r_1,r_2)=\Psi_i^{-\alpha}(p_1-r_1,r_2)=\Psi_i^{-\alpha}(r_1,p_2-r_2)=\Psi_i^{\alpha}(p_1-r_1,p_2-r_2)$ for $i=1,2$ by (3.3.3)-(3.3.6). Now we set

$$B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2)=\frac{1}{\Phi^{\alpha}(r_1,r_2)}b^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2).$$

4.2. **Indecomposable left ideals $P^{\alpha}_{r_1,p_2}(s_1,s_2)$ and $P^{\alpha}_{p_1,r_2}(s_1,s_2)$.** Let us first fix $1\le r_1\le p_1-1$ and $r_2=p_2$. We set

$$\begin{aligned}
(4.2.1)\qquad &B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)\\
=&\frac{e_1^{p_1-r_1-1-k_1}f_2^{n_2}\sum_{m_1=1}^{p_1-r_1}\gamma^{\alpha}_{m_1}(r_1,p_2)e_1^{p_1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,p_2}(s_1,s_2)}{\Phi^{\alpha}(r_1,p_2)\prod_{j_1=k_1+1}^{p_1-r_1-1}\varphi_1^{-\alpha}(j_1,p_1-r_1,p_2)},
\end{aligned}$$

$$
\begin{aligned}
&B^{\alpha,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)\\
=&\frac{f_1^{n_1}f_2^{n_2}}{\Phi^{\alpha}(r_1,p_2)}\sum_{m_1=1}^{p_1-r_1}\gamma^{\alpha}_{m_1}(r_1,p_2)e_1^{p_1-1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^{\alpha}_{r_1,r_2}(r_1,r_2)\\
&\qquad -\Psi^{\alpha}_1(r_1,p_2)B^{\alpha}_{n_1,n_2}(r_1,p_2,s_1,s_2),
\end{aligned}\tag{4.2.2}
$$

$$
B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)=f_1^{r_1+k_1}B^{\alpha,\uparrow}_{0,n_2}(r_1,p_2,s_1,s_2),\tag{4.2.3}
$$

for $0\le n_1\le r_1-1$, $0\le k_1\le p_1-r_1-1$ and $0\le n_2\le p_2-1$.

Lemma 4.4. *For $0\le n_2\le p_2-1$, the following relations hold:*

$$
KB^{\alpha,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)=-\alpha q_1^{p_1-r_1-1-2k_1}q_2^{r_2-1-2n_2}B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2),\tag{4.2.4}
$$

$$
KB^{\alpha,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)=\alpha q_1^{r_1-1-2n_1}q_2^{r_2-1-2n_2}B^{\alpha,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2),\tag{4.2.5}
$$

$$
KB^{\alpha,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)=-\alpha q_1^{p_1-r_1-1-2k_1}q_2^{r_2-1-2n_2}B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2),\tag{4.2.6}
$$

$$
\begin{aligned}
&e_1B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)\\
&=\begin{cases}\varphi_1^{-\alpha}(k_1,p_1-r_1,p_2)B^{\alpha,\leftarrow}_{k_1-1,n_2}(r_1,p_2,s_1,s_2), & 1\le k_1\le p_1-r_1-1,\\ 0, & k_1=0,\end{cases}
\end{aligned}\tag{4.2.7}
$$

$$
\begin{aligned}
&f_1B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)\\
&=\begin{cases}B^{\alpha,\leftarrow}_{k_1+1,n_2}(r_1,p_2,s_1,s_2), & 0\le k_1\le p_1-r_1-2,\\ B^{\alpha,\downarrow}_{0,n_2}(r_1,p_2,s_1,s_2), & k_1=p_1-r_1-1,\end{cases}
\end{aligned}\tag{4.2.8}
$$

$$
f_1B^{\alpha,\downarrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)=\begin{cases}B^{\alpha,\downarrow}_{n_1+1,n_2}(r_1,p_2,s_1,s_2), & 0\le n_1\le r_1-2,\\ 0, & n_1=r_1-1,\end{cases}\tag{4.2.9}
$$

$$
\begin{aligned}
&e_1B^{\alpha,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)\\
&\quad=\begin{cases}\varphi_1^{\alpha}(n_1,r_1,p_2)B^{\alpha,\uparrow}_{n_1-1,n_2}(r_1,p_2,s_1,s_2)+B^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,p_2,s_1,s_2), & 1\le n_1\le r_1-1,\\ B^{\alpha,\uparrow}_{p_1-r_1-1,n_2}(r_1,p_2,s_1,s_2), & n_1=0,\end{cases}
\end{aligned}\tag{4.2.10}
$$

$$
\begin{aligned}
&f_1B^{\alpha,\uparrow}_{n_1-1,n_2}(r_1,p_2,s_1,s_2)\\
&\quad=\begin{cases}B^{\alpha,\uparrow}_{n_1+1,n_2}(r_1,p_2,s_1,s_2), & 0\le n_1\le r_1-2,\\ B^{\alpha,\rightarrow}_{0,n_2}(r_1,p_2,s_1,s_2), & n_1=r_1-1,\end{cases}
\end{aligned}\tag{4.2.11}
$$

$$\begin{aligned}&e_1 B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)\\ &=\begin{cases}\varphi_1^{-\alpha}(k_1,p_1-r_1,p_2)B^{\alpha,\rightarrow}_{k_1-1,n_2}(r_1,p_2,s_1,s_2), & 1\le k_1\le p_1-r_1-1,\\ B^{\alpha,\downarrow}_{r_1-1,n_2}(r_1,p_2,s_1,s_2), & k_1=0,\end{cases}\end{aligned}\tag{4.2.12}$$

$$\begin{aligned}&f_1 B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)\\ &=\begin{cases}B^{\alpha,\rightarrow}_{k_1+1,n_2}(r_1,p_2,s_1,s_2), & 0\le k_1\le p_1-r_1-2,\\ 0, & k_1=p_1-r_1-1.\end{cases}\end{aligned}\tag{4.2.13}$$

Proof. We can see (4.2.4)-(4.2.6), (4.2.7), (4.2.11) and (4.2.13) by the definition.

By Lemma 4.1, Proposition 4.3 and (4.2.1), we obtain (4.2.8). Then we can see $0=f_1^{p_1}B^{\alpha,\leftarrow}_{0,n_2}(r_1,p_2,s_1,s_2)=B^{\alpha,\downarrow}_{r_1,n_2}(r_1,p_2,s_1,s_2)$ by (4.1.2), which proves (4.2.9). By Lemma 4.2 and Proposition 4.3, we have $e_1B^{\alpha,\uparrow}_{0,n_2}(r_1,p_2,s_1,s_2)=B^{\alpha,\leftarrow}_{p_1-r_1-1,n_2}(r_1,p_2,s_1,s_2)$ and then we can see that

$$\begin{aligned}e_1B^{\alpha,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)&=(f_1^{n_1}e_1+[e_1,f_1^{n_1}])B^{\alpha,\uparrow}_{0,n_2}(r_1,p_2,s_1,s_2)\\ &=\varphi_1^{\alpha}(n_1,r_1,p_2)B^{\alpha,\uparrow}_{n_1-1,n_2}(r_1,p_2,s_1,s_2)+B^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,p_2,s_1,s_2)\end{aligned}$$

for $1\le n_1\le r_1-1$ by Lemma 4.1 and (4.2.8). Since $\varphi_1^{\alpha}(r_1+k_1,r_1,p_2)=\varphi_1^{-\alpha}(k_1,p_1-r_1,p_2)$, we have (4.2.12). □

By Lemma 4.1 we have the following.

Lemma 4.5.

$$\begin{aligned}&e_2 B^{\alpha,\bullet}_{n_1,n_2}(r_1,p_2,s_1,s_2)\\ &=\begin{cases}\varphi_2^{\alpha}(n_2,r_1,p_2)B^{\alpha,\bullet}_{n_1,n_2-1}(r_1,p_2,s_1,s_2), & 1\le n_2\le p_2-1,\\ 0, & n_2=0,\end{cases}\end{aligned}\tag{4.2.14}$$

$$\begin{aligned}&f_2 B^{\alpha,\bullet}_{n_1,n_2}(r_1,p_2,s_1,s_2)\\ &=\begin{cases}B^{\alpha,\bullet}_{n_1,n_2+1}(r_1,p_2,s_1,s_2), & 0\le n_2\le p_2-2,\\ 0, & n_2=p_2-1,\end{cases}\end{aligned}\tag{4.2.15}$$

where $0\le n_1\le r_1$ if $\bullet\in\{\uparrow,\downarrow\}$ and $0\le n_1\le p_1-r_1-1$ if $\bullet\in\{\leftarrow,\rightarrow\}$.

Let $X^{\alpha}_{r_1,p_2}(s_1,s_2)$ be the space spanned by the set

$$\{B^{\alpha,\downarrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)|0\le n_1\le r_1-1,0\le n_2\le p_2-1\}.$$

Then $X^{\alpha}_{r_1,p_2}(s_1,s_2)$ is isomorphic to the simple module $X^{\alpha}_{r_1,p_2}$ by (4.1.4), (4.2.9) and (4.2.14)-(4.2.15).

By Lemma 4.2, Lemma 4.4 and Lemma 4.5, we have the following.

Proposition 4.6. *Let $P^{\alpha}_{r_1,p_2}(s_1,s_2)$ be the space spanned by vectors of the form*

$$\begin{aligned}&B^{\alpha,\uparrow\downarrow}_{n_1,n_2}(r_1,p_2,s_1,s_2),0\le n_1\le r_1-1,\ 0\le n_2\le p_2-1,\\ &B^{\alpha,\rightleftarrows}_{k_1,n_2}(r_1,p_2,s_1,s_2),0\le k_1\le p_1-r_1-1,\ 0\le n_2\le p_2-1\end{aligned}$$

for $\alpha=\pm$, $1\le r_1\le p_1-1$, $1\le s_1\le r_1$ and $1\le s_2\le p_2$. Then $P^{\alpha}_{r_1,p_2}(s_1,s_2)$ is a $2p_1p_2$-dimensional indecomposable left ideal whose socle is isomorphic to the simple *module $X^{\alpha}_{r_1,p_2}$. In particular, for fixed $1 \leq r_1 \leq p_1 - 1$, the left ideals $P^{\alpha}_{r_1,p_2}(s_1, s_2)$ for $1 \leq s_1 \leq r_1$ and $1 \leq s_2 \leq p_2$ are isomorphic to each other.*

By Proposition 4.6, we can write $P^{\alpha}_{r_1,p_2} \cong P^{\alpha}_{r_1,p_2}(s_1, s_2)$ for $1 \leq r_1 \leq p_1 - 1$ and $\alpha = \pm$. We denote a basis of $P^{\alpha}_{r_1,p_2}$ by

$$\begin{aligned}&\mathsf{b}^{\alpha,\uparrow\downarrow}_{n_1,n_2}(r_1, p_2),\ 0 \leq n_1 \leq r_1 - 1,\ 0 \leq n_2 \leq p_2 - 1,\\ &\mathsf{b}^{\alpha,\rightleftarrows}_{k_1,n_2}(r_1, p_2),\ 0 \leq k_1 \leq p_1 - r_1 - 1,\ 0 \leq n_2 \leq p_2 - 1.\end{aligned}$$

This basis corresponds to the basis $B^{\alpha,\bullet}$ of $P^{\alpha}_{r_1,p_2}(s_1, s_2)$.

We now fix $r_1 = p_1$, $1 \leq r_2 \leq p_2 - 1$ and set

$$\begin{aligned}&B^{\alpha,\leftarrow}_{n_1,k_2}(p_1, r_2, s_1, s_2)\\ &=\frac{f_1^{n_1} e_2^{p_2-r_2-1-k_2} \sum_{m_2=1}^{p_2-r_2} \delta^{\alpha}_{m_2}(p_1, r_2) e_1^{p_1-1} e_2^{p_2-m_2} f_1^{p_1-s_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{p_1,r_2}(s_1, s_2)}{\Phi^{\alpha}(p_1, r_2) \prod_{j_2=k_2+1}^{p_2-r_2-1} \varphi_2^{-\alpha}(j_2, p_1, p_2 - r_2)},\end{aligned} \tag{4.2.16}$$

$$\begin{aligned}&B^{\alpha,\uparrow}_{n_1,n_2}(p_1, r_2, s_1, s_2)\\ &= \frac{f_1^{n_1} f_2^{n_2}}{\Phi^{\alpha}(p_1, r_2)} \sum_{m_2=1}^{p_2-r_2} \delta^{\alpha}_{m_2}(p_1, r_2) e_1^{p_1-1} e_2^{p_2-1-m_2} f_1^{p_1-s_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{p_1,r_2}(s_1, s_2)\\ &\qquad - \Psi^{\alpha}_2(p_1, r_2) B^{\alpha,\downarrow}_{n_1,n_2}(p_1, r_2, s_1, s_2),\end{aligned} \tag{4.2.17}$$

$$B^{\alpha,\rightarrow}_{n_1,k_2}(p_1, r_2, s_1, s_2) = f_2^{r_2+k_2} B^{\alpha,\uparrow}_{n_1,0}(p_1, r_2, s_1, s_2), \tag{4.2.18}$$

for $0 \leq n_1 \leq p_1 - 1$, $0 \leq n_2 \leq r_2 - 1$ and $0 \leq k_2 \leq p_2 - r_2 - 1$. Then we have the following.

Proposition 4.7. *Let $P^{\alpha}_{p_1,r_2}(s_1, s_2)$ be the space spanned by vectors of the form*

$$\begin{aligned}&B^{\alpha,\uparrow\downarrow}_{n_1,n_2}(p_1, r_2, s_1, s_2), 0 \leq n_1 \leq p_1 - 1,\ 0 \leq n_2 \leq r_2 - 1,\\ &B^{\alpha,\rightleftarrows}_{n_1,k_2}(p_1, r_2, s_1, s_2), 0 \leq n_1 \leq p_1 - 1,\ 0 \leq k_2 \leq p_2 - r_2 - 1\end{aligned}$$

for $\alpha = \pm$, $1 \leq r_2 \leq p_2 - 1$, $1 \leq s_1 \leq p_1$ and $1 \leq s_2 \leq r_2$. Then $P^{\alpha}_{p_1,r_2}(s_1, s_2)$ is a $2p_1p_2$-dimensional indecomposable left ideal whose socle is isomorphic to the simple module $X^{\alpha}_{p_1,r_2}$. In particular, for fixed $1 \leq r_2 \leq p_2 - 1$, the modules $P^{\alpha}_{p_1,r_2}(s_1, s_2)$ for $1 \leq s_1 \leq p_1$ and $1 \leq s_2 \leq r_2$ are isomorphic to each other.

By Proposition 4.7, we can also write $P^{\alpha}_{p_1,r_2} \cong P^{\alpha}_{p_1,r_2}(s_1, s_2)$ for $\alpha = \pm$ and $1 \leq r_2 \leq p_2 - 1$. We denote the basis of $P^{\alpha}_{p_1,r_2}$ which corresponds to the basis $B^{\alpha,\bullet}(s_1, s_2)$ of $P^{\alpha}_{p_1,r_2}(s_1, s_2)$ is denoted by

$$\begin{aligned}&\mathsf{b}^{\alpha,\uparrow\downarrow}_{n_1,n_2}(p_1, r_2),\ 0 \leq n_1 \leq p_1 - 1,\ 0 \leq n_2 \leq r_2 - 1,\\ &\mathsf{b}^{\alpha,\rightleftarrows}_{n_1,k_2}(r_1, p_2),\ 0 \leq k_1 \leq p_1 - 1,\ 0 \leq n_2 \leq p_2 - r_2 - 1,\end{aligned}$$

4.3. **Indecomposable left ideal $P^{\alpha}_{r_1,r_2}$.** Let us fix $1 \le r_1 \le p_1 - 1$ and $1 \le r_2 \le p_2 - 1$. Set

(4.3.1)
$$B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) = \frac{e_1^{p_1-r_1-1-k_1} f_2^{n_2}}{\Phi^{\alpha}(r_1,r_2)\prod_{j_1=k_1+1}^{p_1-r_1-1}\varphi_1^{\alpha}(j_1,p_1-r_1,r_2)} \times \sum_{m_1=1}^{p_1-r_1} \gamma^{\alpha}_{m_1}(r_1,r_2) e_1^{p_1-m_1} e_2^{p_2-1} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2} v^{\alpha}_{r_1,r_2}(s_1,s_2),$$

(4.3.2)
$$\begin{aligned} &B^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) \\ =&\frac{f_1^{n_1} f_2^{n_2}}{\Phi^{\alpha}(r_1,r_2)}\{\sum_{m_1=1}^{p_1-r_1} \gamma^{\alpha}_{m_1}(r_1,r_2) e_1^{p_1-1-m_1} e_2^{p_2-1} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2} v^{\alpha}_{r_1,r_2}(s_1,s_2)\} \\ &- \Psi^{\alpha}_1(r_1,r_2) B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2), \end{aligned}$$

(4.3.3)
$$B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) = f_1^{r_1+k_1} B^{\alpha,\uparrow}_{0,n_2}(r_1,r_2,s_1,s_2),$$

(4.3.4)
$$\begin{aligned} &L^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ =&\frac{f_1^{n_1} e_2^{p_2-r_2-1-k_2}}{\Phi^{\alpha}(r_1,r_2)\prod_{j_2=k_2+1}^{p_2-r_2-1}\varphi_2^{-\alpha}(j_2,r_1,p_2-r_2)} \\ &\times \sum_{m_2=1}^{p_2-r_2} \delta^{\alpha}_{m_2}(r_1,r_2) e_1^{p_1-1} e_2^{p_2-m_2} f_1^{p_1-s_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{r_1,r_2}(s_1,s_2), \end{aligned}$$

(4.3.5)
$$\begin{aligned} &L^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) \\ =&\frac{e_1^{p_1-r_1-1-k_1} e_2^{p_2-r_2-1-k_2}}{\Phi^{\alpha}(r_1,r_2)\prod_{j_1=k_1+1}^{p_1-r_1-1}\prod_{j_2=k_2+1}^{p_2-r_2-1}\varphi_1^{-\alpha}(j_1,p_1-r_1,r_2)\varphi_2^{-\alpha}(j_2,r_1,p_2-r_2)} \\ &\times \sum_{m_1=1}^{p_1-r_1}\sum_{m_2=1}^{p_2-r_2} \gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2) e_1^{p_1-m_1} e_2^{p_2-m_2} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{r_1,r_2}(s_1,s_2), \end{aligned}$$

(4.3.6)
$$\begin{aligned} &L^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ &= \frac{f_1^{n_1} e_2^{p_2-r_2-1-k_2}}{\Phi^{\alpha}(r_1,r_2)\prod_{j_2=k_2+1}^{p_2-r_2-1}\varphi_2 - \alpha(j_2,r_1,p_2-r_2)} \\ &\times \{\sum_{m_1=1}^{p_1-r_1}\sum_{m_2=1}^{p_2-r_2} \gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2) e_1^{p_1-1-m_1} e_2^{p_2-m_2} f_1^{p_1-s_1-m_1} f_2^{p_2-s_2-m_2} v^{\alpha}_{r_1,r_2}(s_1,s_2)\} \\ &- \Psi^{\alpha}_1(r_1,r_2) L^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2), \end{aligned}$$

(4.3.7)
$$L^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) = f_1^{r_1+k_1} L^{\alpha,\uparrow}_{0,k_2}(r_1,r_2,s_1,s_2),$$

(4.3.8)

$$
\begin{aligned}
&T^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2)\\
=&\frac{f_1^{n_1}f_2^{n_2}}{\Phi^\alpha(r_1,r_2)}\{\sum_{m_2=1}^{p_2-r_2}\delta^\alpha_{m_2}(r_1r_2)e^{p_1-1}e_2^{p_2-1-m_2}f_1^{p_1-s_1}f_2^{p_2-s_2-m_2}v^\alpha_{r_1,r_2}(s_1,s_2)\}\\
&\qquad\qquad\qquad - \Psi^\alpha_2(r_1,r_2)B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2),
\end{aligned}
$$

(4.3.9)

$$
\begin{aligned}
&T^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) = \frac{e_1^{p_1-r_1-1-k_1}f_2^{n_2}}{\Phi^\alpha(r_1,r_2)\prod_{j_1=k_1+1}^{p_1-r_1-1}\varphi_1^{-\alpha}(j_1,p_1-r_1,r_2)}\\
&\times\{\sum_{m_1=1}^{p_1-r_1}\sum_{m_2=1}^{p_2-r_2}\gamma^\alpha_{m_1}(r_1,r_2)\delta^\alpha_{m_2}(r_1,r_2)e_1^{p_1-m_1}e_2^{p_2-1-m_2}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2-m_2}v^\alpha_{r_1,r_2}(r_1,r_2)\}\\
&\qquad\qquad - \Psi^\alpha_2(r_1,r_2)B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),
\end{aligned}
$$

(4.3.10)

$$
\begin{aligned}
&T^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) = \frac{f_1^{n_1}f_2^{n_2}}{\Phi^\alpha(r_1,r_2)}\\
&\times\Big\{\sum_{m_1=1}^{p_1-r_1}\sum_{m_2=1}^{p_2-r_2}\gamma^\alpha_{m_1}(r_1,r_2)\delta^\alpha_{m_2}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1-m_2}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2-m_2}v^\alpha_{r_1,r_2}(s_1,s_2)\\
&-\Psi^\alpha_1(r_1,r_2)\sum_{m_2=1}^{p_2-r_2}\delta^\alpha_{m_2}(r_1r_2)e^{p_1-1}e_2^{p_2-1-m_2}f_1^{p_1-s_1}f_2^{p_2-s_2-m_2}v^\alpha_{r_1,r_2}(s_1,s_2)\\
&-\Psi^\alpha_2(r_1,r_2)\sum_{m_1=1}^{p_1-r_1}\gamma^\alpha_{m_1}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}v^\alpha_{r_1,r_2}(s_1,s_2)\\
&+\Psi^\alpha_1(r_1,r_2)\Psi^\alpha_2(r_1,r_2)e_1^{p_1-1}e_2^{p_2-1}f_1^{p_1-s_1}f_2^{p_2-s_2}v^\alpha_{r_1,r_1}(s_1,s_2)\Big\},
\end{aligned}
$$

(4.3.11)

$$T^{\alpha,\rightarrow}_{k_1,r_2}(r_1,r_2,s_1,s_2) = f_1^{r_1+k_1}T^{\alpha,\uparrow}_{0,n_2}(r_1,r_2,s_1,s_2),$$

(4.3.12)

$$R^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) = f_2^{r_2+k_2}T^{\alpha,\downarrow}_{n_1,0}(r_1,r_2,s_1,s_2),$$

(4.3.13)

$$R^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) = f_2^{r_2+k_2}T^{\alpha,\leftarrow}_{k_1,0}(r_1,r_2,s_1,s_2),$$

(4.3.14)

$$R^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) = f_2^{r_2+k_2}T^{\alpha,\uparrow}_{n_1,0}(r_1,r_2,s_1,s_2),$$

(4.3.15)

$$R^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) = f_2^{r_2+k_2}T^{\alpha,\rightarrow}_{k_1,0}(r_1,r_2,s_1,s_2),$$

for $0 \le n_i \le r_i - 1$ and $0 \le k_i \le p_i - r_i - 1$. By Lemma 4.1 and Proposition 4.3, we can see

$$\begin{aligned} &e_1 B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} \varphi_1^{-\alpha}(k_1, p_1 - r_1, r_2) B^{\alpha,\leftarrow}_{k_1-1,n_2}(r_1,r_2,s_1,s_2), & 1 \le k_1 \le p_1 - r_1 - 1 \\ 0, & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.16}$$

$$\begin{aligned} &f_1 B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} B^{\alpha,\leftarrow}_{k_1+1,n_2}(r_1,r_2,s_1,s_2), & 0 \le k_1 \le p_1 - r_1 - 2, \\ B^{\alpha,\downarrow}_{0,n_2}(r_1,r_2,s_1,s_2), & k_1 = p_1 - r_1 - 1. \end{cases} \end{aligned} \tag{4.3.17}$$

Thus we have $B^{\alpha,\downarrow}_{r_1,n_2}(r_1,r_2,s_1,s_2) = f_1^{p_1} B^{\alpha,\leftarrow}_{0,n_2}(r_1,r_2,s_1,s_2) = 0$ for $0 \le n_2 \le r_2 - 1$, which shows

$$f_1 B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) = \begin{cases} B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2), & 0 \le n_1 \le r_1 - 2, \\ 0, & n_1 = r_1 - 1. \end{cases} \tag{4.3.18}$$

By Lemma 4.1, Proposition 4.3 and (4.3.18), we have

(4.3.19)

$$\begin{aligned} &e_1 B^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} \varphi_1^{\alpha}(n_1, r_1, r_2) B^{\alpha,\uparrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2) + B^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2), & 1 \le n_1 \le r_1 - 1, \\ B^{\alpha,\leftarrow}_{p_1-r_1-1,n_2}(r_1,r_2,s_1,s_2), & n_1 = 0, \end{cases} \end{aligned}$$

(4.3.20)

$$\begin{aligned} &f_1 B^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} B^{\alpha,\uparrow}_{n_1+1,n_2}(r_1,r_2,s_1,s_2), & 0 \le n_1 \le r_1 - 2, \\ B^{\alpha,\rightarrow}_{0,n_2}(r_1,r_2,s_1,s_2), & n_1 = r_1 - 1, \end{cases} \end{aligned}$$

(4.3.21)

$$\begin{aligned} &e_1 B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} \varphi_1^{-\alpha}(k_1, p_1 - r_1, r_2) B^{\alpha,\rightarrow}_{k_1-1,n_2}(r_1,r_2,s_1,s_2), & 1 \le k_1 \le p_1 - r_1 - 1 \\ B^{\alpha,\downarrow}_{r_1-1,n_2}(r_1,r_2,s_1,s_2), & k_1 = 0, \end{cases} \end{aligned}$$

(4.3.22)

$$\begin{aligned} &f_1 B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} B^{\alpha,\rightarrow}_{k_1+1,n_2}(r_1,r_2,s_1,s_2), & 0 \le k_1 \le p_1 - r_1 - 2, \\ 0, & k_1 = p_1 - r_1 - 1, \end{cases} \end{aligned}$$

and we also have

$$\begin{aligned} &e_2 B^{\alpha,\bullet}_{n_1,n_2}(r_1,r_2,s_1,s_2) \\ &= \begin{cases} \varphi_2^{\alpha}(n_2, r_1, r_2) B^{\alpha,\bullet}_{n_1,n_2-1}(r_1,r_2,s_1,s_2), & 1 \le n_2 \le r_2 - 1, \\ 0, & n_2 = 0, \end{cases} \end{aligned} \tag{4.3.23}$$

where $0 \leq n_1 \leq r_1 - 1$ if $\bullet \in \{\uparrow, \downarrow\}$ and $0 \leq n_1 \leq p_1 - r_1 - 1$ if $\bullet \in \{\leftarrow, \rightarrow\}$. By Lemma 4.1 and (4.3.4)-(4.3.7), we obtain

$$\begin{aligned} &e_1 L^{\alpha,\downarrow}_{n_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} \varphi^{\alpha}_1(n_1, r_1, r_2) L^{\alpha,\downarrow}_{n_1-1,k_2}(r_1, r_2, s_1, s_2), & 1 \leq n_1 \leq r_1 - 1, \\ 0, & n_1 = 0, \end{cases} \end{aligned} \tag{4.3.24}$$

$$\begin{aligned} &e_1 L^{\alpha,\leftarrow}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} \varphi^{\alpha}_1(k_1, r_1, p_2 - r_2) L^{\alpha,\leftarrow}_{k_1-1,k_2}(r_1, r_2, s_1, s_2), & 1 \leq k_1 \leq p_1 - r_1 - 1, \\ 0, & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.25}$$

$$\begin{aligned} &f_1 L^{\alpha,\leftarrow}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} L^{\alpha,\leftarrow}_{k_1+1,k_2}(r_1, r_2, s_1, s_2), & 0 \leq k_1 \leq p_1 - r_1 - 2, \\ L^{\alpha,\downarrow}_{0,k_2}(r_1, r_2, s_1, s_2), & k_1 = p_1 - r_1 - 1, \end{cases} \end{aligned} \tag{4.3.26}$$

where the last relation follows from Proposition 4.3. By (4.3.4) and (4.3.26), we see $L^{\alpha,\downarrow}_{r_1,k_2}(r_1, r_2, s_1, s_2) = f_1^{p_1} L^{\alpha,\leftarrow}_{0,k_2}(r_1, r_2, s_1, s_2) = 0$. Hence we have

$$\begin{aligned} &f_1 L^{\alpha,\downarrow}_{n_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} L^{\alpha,\downarrow}_{n_1+1,k_2}(r_1, r_2, s_1, s_2), & 0 \leq n_1 \leq r_1 - 2, \\ 0, & n_1 = r_1 - 1. \end{cases} \end{aligned} \tag{4.3.27}$$

We can see that

$$\begin{aligned} &e_1 L^{\alpha,\uparrow}_{n_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} \varphi^{\alpha}_1(n_1, r_1, r_2) L^{\alpha,\uparrow}_{n_1-1,k_2}(r_1, r_2, s_1, s_2) + L^{\alpha,\downarrow}_{n_1-1,k_2}(r_1, r_2, s_1, s_2), & 1 \leq n_1 \leq r_1 - 1, \\ L^{\alpha,\leftarrow}_{p_1-r_1-1,k_2}(r_1, r_2, s_1, s_2), & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.28}$$

$$f_1 L^{\alpha,\uparrow}_{n_1,k_2}(r_1, r_2, s_1, s_2) = \begin{cases} L^{\alpha,\uparrow}_{n_1+1,k_2}(r_1, r_2, s_1, s_2), & 0 \leq n_1 \leq r_1 - 2, \\ L^{\alpha,\rightarrow}_{0,k_2}(r_1, r_2, s_1, s_2), & n_1 = p_1 - r_1 - 1. \end{cases} \tag{4.3.29}$$

by Lemma 4.1 and we also have

$$\begin{aligned} &e_1 L^{\alpha,\rightarrow}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} \varphi^{-\alpha}_1(k_1, p_1 - r_1, r_2) L^{\alpha,\uparrow}_{k_1-1,k_2}(r_1, r_2, s_1, s_2), & 1 \leq k_1 \leq p_1 - r_1 - 1, \\ L^{\alpha,\downarrow}_{0,k_2}(r_1, r_2, s_1, s_2), & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.30}$$

$$\begin{aligned} &f_1 L^{\alpha,\rightarrow}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\ &= \begin{cases} L^{\alpha,\rightarrow}_{k_1+1,k_2}(r_1, r_2, s_1, s_2), & 0 \leq k_1 \leq p_1 - r_1 - 2, \\ 0, & k_1 = p_1 - r_1 - 1, \end{cases} \end{aligned} \tag{4.3.31}$$

by Lemma 4.1 and Proposition 4.3, in addition, the last relation follows from (4.3.7) and $f_1^{p_1} = 0$. Moreover, we have

$$\begin{aligned}
&e_2 L^{\alpha,\bullet}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} \varphi_2^{-\alpha}(r_1, p_2 - r_2) L^{\alpha,\bullet}_{k_1,k_2-1}(r_1, r_2, s_1, s_2), & 1 \le k_2 \le p_2 - r_2 - 1, \\ 0, & k_2 = 0, \end{cases}
\end{aligned} \tag{4.3.32}$$

$$\begin{aligned}
&f_2 L^{\alpha,\bullet}_{k_1,k_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} L^{\alpha,\bullet}_{k_1,k_2+1}(r_1, r_2, s_1, s_2), & 0 \le k_2 \le p_2 - r_2 - 2, \\ B^{\alpha,\bullet}_{k_1,0}(r_1, r_2, s_1, s_2), & k_2 = p_2 - r_2 - 1, \end{cases}
\end{aligned} \tag{4.3.33}$$

where $0 \le k_1 \le r_1 - 1$ if $\bullet \in \{\uparrow, \downarrow\}$ and $0 \le k_1 \le p_1 - r_1 - 1$ if $\bullet \in \{\leftarrow, \rightarrow\}$. By (4.3.33), we can see $b^{\alpha,\bullet}_{k_1,r_2}(r_1, r_2, s_1, s_2) = f_2^{p_2} l^{\alpha,\bullet}_{k_1,0}(r_1, r_2, s_1, s_2) = 0$. Therefore we have

$$f_2 B^{\alpha,\bullet}_{n_1,n_2}(r_1, r_2, s_1, s_2) = \begin{cases} B^{\alpha,\bullet}_{n_1,n_2+1}(r_1, r_2, s_1, s_2), & 0 \le n_2 \le r_1 - 2, \\ 0, & n_2 = r_2 - 1, \end{cases} \tag{4.3.34}$$

where $0 \le n_1 \le r_1 - 1$ if $\bullet \in \{\uparrow, \downarrow\}$ and $0 \le n_1 \le p_1 - r_1 - 1$ if $\bullet \in \{\leftarrow, \rightarrow\}$. The following relations are obtained by similar argument used to show (4.3.16)-(4.3.22):

$$\begin{aligned}
&T^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} \varphi_1^{\alpha}(n_1, r_1, r_2) T^{\alpha,\downarrow}_{n_1-1,n_2}(r_1, r_2, s_1, s_2), & 1 \le n_1 \le r_1 - 1, \\ 0, & n_1 = 0, \end{cases}
\end{aligned} \tag{4.3.35}$$

$$\begin{aligned}
&f_1 T^{\alpha,\downarrow}_{n_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} T^{\alpha,\downarrow}_{n_1+1,n_2}(r_1, r_2, s_1, s_2), & 0 \le n_1 \le r_1 - 2, \\ 0, & n_1 = r_1 - 1, \end{cases}
\end{aligned} \tag{4.3.36}$$

$$\begin{aligned}
&e_1 T^{\alpha,\leftarrow}_{k_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} \varphi_1^{-\alpha}(k_1, p_1 - r_1, r_2) T^{\alpha,\downarrow}_{k_1-1,n_2}(r_1, r_2, s_1, s_2), & 1 \le k_1 \le p_1 - r_1 - 1, \\ 0, & k_1 = 0, \end{cases}
\end{aligned} \tag{4.3.37}$$

$$\begin{aligned}
&f_1 T^{\alpha,\leftarrow}_{k_1,n_2}(r_1, r_2, s_1, s_2) \\
&= \begin{cases} T^{\alpha,\leftarrow}_{k_1+1,n_2}(r_1, r_2, s_1, s_2), & 0 \le k_1 \le p_1 - r_1 - 2, \\ T^{\alpha,\downarrow}_{0,n_2}(r_1, r_2, s_1, s_2), & k_1 = p_1 - r_1 - 1, \end{cases}
\end{aligned} \tag{4.3.38}$$

$$e_1T^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) \tag{4.3.39}$$
$$=\begin{cases}\varphi^{\alpha}_1(n_1,r_1,r_2)T^{\alpha,\uparrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2)+T^{\alpha,\downarrow}_{n_1-1,n_2}(r_1,r_2,s_1,s_2), & 1\le n_1\le r_1-1,\\ T^{\alpha,\leftarrow}_{p_1-r_1-1,n_2}(r_1,r_2,s_1,s_2), & n_1=0,\end{cases}$$

$$f_1T^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2) \tag{4.3.40}$$
$$=\begin{cases}T^{\alpha,\uparrow}_{n_1+1,n_2}(r_1,r_2,s_1,s_2), & 0\le n_1\le r_1-2,\\ T^{\alpha,\rightarrow}_{0,n_2}(r_1,r_2,s_1,s_2), & n_1=r_1-1,\end{cases}$$

$$e_1T^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \tag{4.3.41}$$
$$=\begin{cases}\varphi^{-\alpha}_1(k_1,p_1-r_1,r_2)T^{\alpha,\rightarrow}_{k_1-1,n_2}(r_1,r_2,s_1,s_2), & 1\le k_1\le p_1-r_1-1,\\ T^{\alpha,\downarrow}_{r_1-1,n_2}(s_1,s_2), & k_1=0,\end{cases}$$

$$f_1T^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2) \tag{4.3.42}$$
$$=\begin{cases}T^{\alpha,\rightarrow}_{k_1+1,n_2}(r_1,r_2,s_1,s_2), & 0\le k_1\le p_1-r_1-2,\\ 0, & k_1=p_1-r_1-1.\end{cases}$$

By (4.3.8)-(4.3.15), Lemma 4.1 and Proposition 4.3, we obtain

(4.3.43)

$$e_2T^{\alpha,\bullet}_{n_1,n_2}(s_1,s_2)$$
$$=\begin{cases}\varphi^{\alpha}_2(n_2,r_1,r_2)T^{\alpha,\bullet}_{n_1,n_2-1}(r_1,r_2,s_1,s_2)+B^{\alpha,\bullet}_{n_1-1,n_2}(r_1,r_2,s_1,s_2), & 1\le n_2\le r_2-1,\\ L^{\alpha,\bullet}_{n_1,p_2-r_2-1}(r_1,r_2,s_1,s_2), & n_2=0,\end{cases}$$

(4.3.44)

$$f_2T^{\alpha,\bullet}_{n_1,n_2}(r_1,r_2,s_1,s_2)=\begin{cases}T^{\alpha,\bullet}_{n_1,n_2+1}(r_1,r_2,s_1,s_2), & 0\le n_2\le r_2-2,\\ R^{\alpha,\bullet}_{n_1,0}(r_1,r_2,s_1,s_2), & n_2=r_2-1,\end{cases}$$

where $0\le n_1\le r_1-1$ if $\bullet\in\{\uparrow,\downarrow\}$ and $0\le n_1\le p_1-r_1-1$ if $\bullet\in\{\leftarrow,\rightarrow\}$. By (4.3.12)-(4.3.15) and (4.3.35)-(4.3.42), we can see that

$$e_1R^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \tag{4.3.45}$$
$$=\begin{cases}\varphi^{\alpha}_1(n_1,r_1,r_2)R^{\alpha,\uparrow}_{n_1-1,k_2}(r_1,r_2,s_1,s_2)+R^{\alpha,\downarrow}_{n_1-1,k_2}(r_1,r_2,s_1,s_2), & 1\le n_1\le r_1-1,\\ R^{\alpha,\leftarrow}_{p_1-r_1-1,k_2}(r_1,r_2,s_1,s_2), & n_1=0,\end{cases}$$

$$f_1R^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \tag{4.3.46}$$
$$=\begin{cases}R^{\alpha,\uparrow}_{n_1+1,k_2}(r_1,r_2,s_1,s_2), & 0\le n_1\le r_1-2,\\ R^{\alpha,\rightarrow}_{0,k_2}(r_1,r_2,s_1,s_2), & n_1=r_1-1,\end{cases}$$

$$\begin{aligned} & e_1 R^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} \varphi_1^{-\alpha}(k_1,p_1-r_1,r_2) R^{\alpha,\leftarrow}_{k_1-1,k_2}(r_1,r_2,s_1,s_2), & 1 \le k_1 \le p_1-r_1-1, \\ 0, & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.47}$$

$$\begin{aligned} & f_1 R^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} R^{\alpha,\leftarrow}_{k_1+1,k_2}(r_1,r_2,s_1,s_2), & 0 \le k_1 \le p_1-r_1-2, \\ R^{\alpha,\downarrow}_{0,k_2}(r_1,r_2,s_1,s_2), & k_1 = p_1-r_1-1, \end{cases} \end{aligned} \tag{4.3.48}$$

$$\begin{aligned} & e_1 R^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} \varphi_1^{-\alpha}(k_1,p_1-r_1,r_2) R^{\alpha,\rightarrow}_{k_1-1,k_2}(r_1,r_2,s_1,s_2), & 1 \le k_1 \le p_1-r_1-1, \\ R^{\alpha,\downarrow}_{r_1-1,k_2}(r_1,r_2,s_1,s_2), & k_1 = 0, \end{cases} \end{aligned} \tag{4.3.49}$$

$$\begin{aligned} & f_1 R^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} R^{\alpha,\rightarrow}_{k_1+1,k_2}(r_1,r_2,s_1,s_2), & 0 \le k_1 \le p_1-r_1-2, \\ 0, & k_1 = p_1-r_1-1, \end{cases} \end{aligned} \tag{4.3.50}$$

$$\begin{aligned} & e_1 R^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} \varphi_1^{\alpha}(n_1,r_1,r_2) R^{\alpha,\downarrow}_{n_1-1,k_2}(r_1,r_2,s_1,s_2), & 1 \le n_1 \le r_1-1, \\ 0, & n_1 = 0, \end{cases} \end{aligned} \tag{4.3.51}$$

$$\begin{aligned} & f_1 R^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} R^{\alpha,\downarrow}_{n_1+1,k_2}(r_1,r_2,s_1,s_2), & 0 \le n_1 \le r_1-2, \\ 0, & n_1 = 0. \end{cases} \end{aligned} \tag{4.3.52}$$

By Lemma 4.1, (4.3.12)-(4.3.15) and (4.3.35)-(4.3.42), one has

$$\begin{aligned} & e_2 R^{\alpha,\bullet}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} \varphi_2^{-\alpha}(k_2,r_1,p_2-r_2) R^{\alpha,\bullet}_{n_1,k_2-1}(r_1,r_2,s_1,s_2), & 1 \le k_2 \le p_2-r_2-1, \\ B^{\alpha,\bullet}_{n_1,r_2-1}(r_1,r_2,s_1,s_2), & k_2 = 0, \end{cases} \end{aligned} \tag{4.3.53}$$

$$\begin{aligned} & f_2 R^{\alpha,\bullet}_{n_1,k_2}(r_1,r_2,s_1,s_2) \\ & = \begin{cases} R^{\alpha,\bullet}_{n_1,k_2+1}(r_1,r_2,s_1,s_2), & 0 \le k_2 \le p_2-r_2-2, \\ 0, & k_2 = p_2-r_2-1, \end{cases} \end{aligned} \tag{4.3.54}$$

where $0 \le n_1 \le r_1 - 1$ if $\bullet \in \{\uparrow, \downarrow\}$ and $0 \le n_1 \le p_1 - r_1 - 1$ if $\bullet \in \{\leftarrow, \rightarrow\}$.

Let $X^{\alpha}_{r_1,r_2}(s_1,s_2)$ be the space spanned by the vectors

$$\left\{ B^{\alpha}_{n_1,n_2}(r_1,r_2,s_1,s_2); 1 \le n_i \le r_i - 1 \right\}.$$

By (4.1.1)-(4.1.5), (4.3.18) and (4.3.34), $X^{\alpha}_{r_1,r_2}(s_1,s_2)$ is a left module and is isomorphic to the simple module $X^{\alpha}_{r_1,r_2}$.

Proposition 4.8. *Let* $P^{\alpha}_{r_1,r_2}(s_1,s_2)$ *be the space spanned by the vectors*

$$\begin{gathered}
T^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2,s_1,s_2),\ T^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),\ T^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),\ T^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2),\\
L^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2),\ L^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2),\ L^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2),\ L^{\alpha,\downarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),\\
R^{\alpha,\uparrow}_{n_1,k_2}(r_1,r_2,s_1,s_2),\ R^{\alpha,\leftarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2),\ R^{\alpha,\rightarrow}_{k_1,k_2}(r_1,r_2,s_1,s_2),\ R^{\alpha,\downarrow}_{n_1,k_2}(r_1,r_2,s_1,s_2),\\
B^{\alpha,\uparrow}_{n_1,n_2}(r_1,r_2 s_1,s_2),\ B^{\alpha,\leftarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),\ B^{\alpha,\rightarrow}_{k_1,n_2}(r_1,r_2,s_1,s_2),\ B^{\alpha,\downarrow}_{n_1,n_2}(r_1,r_2,s_1,s_2),
\end{gathered}$$

for $0 \le n_i \le r_i - 1$ *and* $0 \le k_i \le p_i - r_i - 1$. *Then* $P^{\alpha}_{r_1,r_2}(s_1,s_2)$ *is a* $4p_1p_2$*-dimensional indecomposable left ideal whose socle is isomorphic to the simple module* $X^{\alpha}_{r_1,r_2}$*, in particular, for fixed* $1 \le r_i \le p_i - 1$ *and* $\alpha = \pm$*, the modules* $P^{\alpha}_{r_1,r_2}(s_1,s_2)$ *for* $1 \le s_i \le r_i$ *are isomorphic to each other.*

By Proposition 4.8, we can write $P^{\alpha}_{r_1,r_2} \cong P^{\alpha}_{r_1,r_2}(s_1,s_2)$. A basis of $P^{\alpha}_{r_1,r_2}$ which corresponds to the basis $T^{\alpha,\bullet}(s_1,s_2), L^{\alpha,\bullet}(s_1,s_2), R^{\alpha,\bullet}(s_1,s_2), B^{\alpha,\bullet}(s_1,s_2)$ is denoted by

$$\mathsf{t}^{\alpha,\bullet}(r_1,r_2), \mathsf{l}^{\alpha,\bullet}(r_1,r_2), \mathsf{r}^{\alpha,\bullet}(r_1,r_2), \mathsf{b}^{\alpha,\bullet}(r_1,r_2).$$

5. PROJECTIVE MODULES AND PRIMITIVE IDEMPOTENTS

In this section we will construct primitive idempotents of $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$ and show that the simple module $X^{\alpha}_{p_1,p_2}$ and indecomposable modules constructed in Section 3 are projective. Moreover, we give the block decomposition of $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$.

5.1. **Primitive idempotents.**

In order to calculate the products of elements in $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$, the following lemma is useful.

Lemma 5.1. (1) *Let* w *be an element in* $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$ *with* $Kw = \alpha q_1^{r_1-1-2n_1} q_2^{r_2-1-2n_2} w$ *for* $0 \le n_1 \le r_1 - 1$ *and* $0 \le n_2 \le r_2 - 1$. *Then*

$$v^{\alpha}_{r_1,r_2}(s_1,s_2)w = \begin{cases} 2p_1p_2 w, & (n_1,n_2) = (s_1-1, s_2-1), \\ 0, & \textit{otherwise}. \end{cases}$$

(2) *Let* w *be an element in* $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$ *with* $Kw = -\alpha q_1^{p_1-r_1-1-2k_1} q_2^{r_2-1-2n_2} w$ *for* $0 \le k_1 \le p_1 - r_1 - 1$ *and* $0 \le n_2 \le r_2 - 1$. *Then* $v^{\alpha}_{r_1,r_2}(s_1,s_2)w = 0$.

(3) *Let* w *be an element in* $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$ *with* $Kw = -\alpha q_1^{r_1-1-2n_1} q_2^{p_2-r_2-1-2k_2} w$ *for* $0 \le n_1 \le r_1 - 1$ *and* $0 \le k_2 \le p_2 - r_2 - 1$. *Then* $v^{\alpha}_{r_1,r_2}(s_1,s_2)w = 0$.

(4) *Let* w *be an element in* $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$ *with* $Kw = \alpha q_1^{p_1-r_1-1-2k_1} q_2^{p_2-r_2-1-2k_2} w$ *for* $0 \le k_1 \le p_1 - r_1 - 1$ *and* $0 \le k_2 \le p_2 - r_2 - 1$. *Then* $v^{\alpha}_{r_1,r_2}(s_1,s_2)w = 0$.

Proof. We will show (1) and (2). The proofs of (3) and (4) are similar to the proof of (2).

(1) We can see

$$\begin{aligned}
v^{\alpha}_{r_1,r_2}(s_1,s_2)w &= \sum_{\ell=0}^{2p_1p_2} (\alpha q_1^{-(r_1-2s_1+1)} q_2^{-(r_2-2s_2+1)})^{\ell} (\alpha q_1^{r_1-1-2n_1} q_2^{r_2-2n_2-2}) w \\
&= \sum_{\ell=0}^{2p_1p_2-1} (q_1^{2(s_1-n_1-1)} q_2^{2(s_2-n_2-1)})^{\ell} w \\
&= \sum_{\ell=0}^{2p_1p_2-1} (q^{4\{p_2(s_1-n_1-1)+p_1(s_2-n_2-1)\}})^{\ell} w
\end{aligned}$$

Since $q = \exp(\sqrt{-1}\pi/2p_1p_2)$, the sum does not vanish if and only if

$$p_2(s_1 - n_1 - 1) + p_1(s_2 - n_2 - 1) = mp_1p_2$$

for some $m \in \mathbb{Z}$. Since

$$n_1 = s_1 - 1 + \frac{p_1}{p_2}(s_2 - n_2 - 1) - mp_1p_2 \tag{5.1.1}$$

and p_1, p_2 are coprime, the right hand side of (5.1.1) does not belong to $\{0, 1, \ldots, r_1 - 1\}$ if $s_2 - n_2 - 1 \neq 0$ and $m \neq 0$. Thus we have $(n_1, n_2) = (s_1 - 1, s_2 - 1)$.

(2) We have

$$\begin{aligned} v^{\alpha}_{r_1,r_2}(s_1, s_2)w &= \sum_{\ell=0}^{2p_1p_2-1} (\alpha q_1^{-(r_1-2s_1+1)} q_2^{-(r_2-2s_2+1)})^{\ell} (-\alpha q_1^{p_1-r_1-1-2k_1} q_2^{r_2-2n_2-2})w \\ &= \sum_{\ell=0}^{2p_1p_2-1} q^{-4\ell(p_2(r_1+k_1-s_1+1)-p_1(s_2-n_2-1))}w. \end{aligned}$$

Then the sum does not vanish if and only if $p_2(r_1 + k_1 - s_1 + 1) - p_1(s_2 - n_2 - 1) = mp_1p_2$ for some $m \in \mathbb{Z}$. Since $1 \leq k_1 + r_1 - s_1 + 1 \leq p_1 - 1$ and $-(r_2 - 1) \leq s_2 - n_2 - 1 \leq r_2 - 1$, we have the assertion. □

Let $X^{\alpha}_{p_1,p_2}(s_1, s_2)$ be the space spanned by the vectors

$$\left\{B^{\alpha,\downarrow}_{n_1,n_2}(p_1, p_2, s_1, s_2); 0 \leq n_i \leq p_i - 1,\ i = 1, 2\right\}.$$

We can see that $X^{\alpha}_{p_1,p_2}(s_1, s_2)$ is isomorphic to the simple module $X^{\alpha}_{p_1,p_2}$.

Proposition 5.2. *The element $B^{\alpha,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2)$ is a primitive idempotent of $\mathfrak{g}_{p_1,p_2}$ for $\alpha = \pm$ and $1 \leq s_i \leq p_i - 1$. In particular, the simple module $X^{\alpha}_{p_1,p_2}$ is projective.*

Proof. By the action of $\mathfrak{g}_{p_1,p_2}$ on simple modules and Lemma 5.1, we have

$$\begin{aligned} &(B^{\alpha,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2))^2 \\ =&\frac{2p_1p_2 f_1^{s_1-1} f_2^{s_2-1}}{\Phi^{\alpha}(p_1, p_2)} e_1^{p_1-1} e_2^{p_2-1} f_1^{p_1-s_1} f_2^{p_2-s_2} B^{\alpha,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2) \\ =&\frac{2p_1p_2}{\Phi^{\alpha}(p_1, p_2)} \prod_{i_1=1}^{p_1-1} \prod_{i_2=1}^{p_2-1} \varphi_1^{\alpha}(i_1, p_1, p_2)\varphi_2^{\alpha}(i_2, p_1, p_2) B^{\alpha,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2) \\ =&B^{\alpha,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2). \end{aligned}$$

□

By similar methods used in [1], we have the following.

Proposition 5.3. *The elements $B^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1, p_2, s_1, s_2)$ and $B^{\alpha,\uparrow}_{s_1-1,s_2-1}(p_1, r_2, s_1, s_2)$ are primitive idempotents. In particular, indecomposable modules $P^{\alpha}_{r_1,p_2}$ and $P^{\alpha}_{p_1,r_2}$ are projective.*

We also have the following.

Proposition 5.4. *The elements*

$$T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2),\ 1\le s_1\le p_1,\ 1\le s_2\le p_2, \tag{5.1.2}$$

are primitive idempotents of $\mathfrak{g}_{\boldsymbol{p_1,p_2}}$. *In particular, the indecomposable module* $P^{\alpha}_{r_1,r_2}$ *is a projective module.*

Proof. It is clear that $T^{\alpha,\uparrow}_{s_1-1,s_2-2}(r_1,r_2,s_1,s_2)$ generate the left ideal $P^{\alpha}_{r_1,r_2}(s_1,s_2)$. By Lemma 5.1, we can see that

$$2p_1p_2e_1^{p_1-1}e_2^{p_2-1}f_1^{p_1-s_1}f_2^{p_2-s_2}T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)=\Phi^{\alpha}(r_1,r_2)B^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2).$$

We also have

$$\begin{aligned}
&2p_1p_2\sum_{m_2=1}^{p_2-r_2}\delta^{\alpha}_{m_2}(r_1r_2)e^{p_1-1}e_2^{p_2-1-m_2}f_1^{p_1-s_1}f_2^{p_2-s_2-m_2}T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&=2p_1p_2\Big\{\delta^{\alpha}_{p_2-r_2}(r_1,r_2)e_1^{p_1-1}e_2^{r_2-1}T^{\alpha,\rightarrow}_{p_1-r_1-1,r_2-1}(r_1,r_2,s_1,s_2)\\
&+\sum_{m_2=1}^{p_2-r_2-1}\delta^{\alpha}_{m_2}(r_1,r_2)R^{\alpha,\rightarrow}_{p_1-r_1-1,p_2-r_2-1-m_2}(r_1,r_2,s_1,s_2)\Big\}\\
&=\Phi^{\alpha}(r_1,r_2)\Big\{T^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)+\sum_{i_2=1}^{r_2-1}\frac{1}{\varphi_2^{\alpha}(i_2,r_1,r_1)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&+\sum_{k_2=2}^{p_2-r_2-1}\frac{1}{\varphi_2^{-\alpha}(k_2,r_1,p_2-r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\Big\}
\end{aligned}$$

and

$$\begin{aligned}
&2p_1p_2\sum_{m_1=1}^{p_1-r_1}\gamma^{\alpha}_{m_1}(r_1r_2)e^{p_1-1-m_1}e_2^{p_2-1}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2}T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&=2p_1p_2\Big\{\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)e_1^{r_1-1}e_2^{p_2-1}R^{\alpha,\uparrow}_{r_1-1,p_2-r_2-1}(r_1,r_2,s_1,s_2)\\
&+\sum_{m_1=1}^{p_1-r_1-1}\gamma^{\alpha}_{m_1}(r_1,r_2)R^{\alpha,\rightarrow}_{p_1-r_1-1-m_1,p_2-r_2-1}(r_1,r_2,s_1,s_2)\Big\}\\
&=\Phi^{\alpha}(r_1,r_2)\Big\{B^{\alpha,\uparrow}_{0,0}(r_1,r_2,s_1,s_2)+\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_1)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&+\sum_{k_1=1}^{p_1-r_1-1}\frac{1}{\varphi_1^{-\alpha}(k_1,p_1-r_1,r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\Big\}.
\end{aligned}$$

By Lemma 5.1 and (4.3.2), we have

$$\begin{aligned}
&2p_1p_2\left\{\sum_{m_1=1}^{p_1-r_1}\sum_{m_2=1}^{p_2-r_2}\gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1-m_2}f_1^{p_1-s_1-m_1}f_2^{p_2-s_2-m_2}\right\}\\
&\qquad\qquad\times T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&=2p_1p_2\Big\{\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)\delta^{\alpha}_{p_2-r_2}(r_1,r_2)e_1^{r_1-1}e_2^{r_2-1}T^{\alpha,\uparrow}_{r_1-1,r_2-1}(r_1,r_2,s_1,s_2)\\
&+\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)\sum_{m_2=1}^{p_2-r_2-1}\delta^{\alpha}_{m_2}(r_1,r_2)e_1^{r_1-1}e_2^{p_2-1-m_2}R^{\alpha,\uparrow}_{r_1-1,p_2-r_2-1-m_2}(r_1,r_2,s_1,s_2)\\
&+\delta^{\alpha}_{p_2-r_2}(r_1,r_2)\sum_{m_1=1}^{p_1-r_1-1}\gamma^{\alpha}_{m_1}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{r_2-1}T^{\alpha,\rightarrow}_{p_1-r_1-1-m_1,r_2-1}(r_1,r_2,s_1,s_2)\\
&\sum_{m_1=1}^{p_1-r_1-1}\sum_{m_2=1}^{p_2-r_2-1}\gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1-m_2}\\
&\qquad\qquad\qquad\times R^{\alpha,\rightarrow}_{p_1-r_1-1-m_1,p_2-r_2-1-m_2}(r_1,r_2,s_1,s_2)\Big\}.
\end{aligned}$$

Then we have

$$\begin{aligned}
&2p_1p_2\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)\delta^{\alpha}_{p_2-r_2}(r_1,r_2)e_1^{r_1-1}e_2^{r_2-1}T^{\alpha,\uparrow}_{r_1-1,r_2-1}(r_1,r_2,s_1,s_2)\\
&=2p_1p_2\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)\delta^{\alpha}_{p_2-r_2}(r_1,r_2)e_2^{r_2-1}\Big\{\prod_{i_1=1}^{r_1-1}\varphi_1^{\alpha}(i_1,r_1,r_2)T^{\alpha,\uparrow}_{0,r_2-1}(r_1,r_2,s_1,s_2)\\
&\qquad+\sum_{i_1=1}^{r_1-1}\prod_{\substack{j_1=1\\ j_1\neq i_1}}^{r_1-1}\varphi_1^{\alpha}(j_1,r_1,r_2)T^{\alpha,\downarrow}_{0,r_2-1}(r_1,r_2,s_1,s_2)\Big\}\\
&=\Phi^{\alpha}(r_1,r_2)\Big\{T^{\alpha,\uparrow}_{0,0}(r_1,r_2,s_1,s_2)+\sum_{i_2=1}^{r_2-1}\frac{1}{\varphi_2^{\alpha}(i_2,r_1,r_2)}B^{\alpha,\uparrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&+\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_2)}T^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&+\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_2)}\sum_{i_2=1}^{r_2-1}\frac{1}{\varphi_2^{\alpha}(j_2,r_1,r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\Big\},
\end{aligned}$$

and

$$\begin{aligned}
&2p_1p_2\gamma^{\alpha}_{p_1-r_1}(r_1,r_2)\sum_{m_2=1}^{p_2-r_2-1}\delta^{\alpha}_{m_2}(r_1,r_2)e_1^{r_1-1}e_2^{p_2-1-m_2}R^{\alpha,\uparrow}_{r_1-1,p_2-r_2-1-m_2}(r_1,r_2,s_1,s_2)\\
&=\Phi^{\alpha}(r_1,r_2)\Big\{\sum_{k_2=1}^{p_2-r_2-1}\frac{1}{\varphi_2^{-\alpha}(k_2,r_1,p_2-r_2)}B^{\alpha,\uparrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&\qquad+\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_2)}\sum_{k_2=1}^{p_2-r_2-1}\frac{1}{\varphi_2^{-\alpha}(k_2,r_1,p_2-r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2).
\end{aligned}$$

Similarly we have

$$
\begin{aligned}
&2p_1p_2\delta^{\alpha}_{p_2-r_2}(r_1,r_2)\sum_{m_1=1}^{p_1-r_1-1}\gamma^{\alpha}_{m_1}(r_1,r_2)e_1^{p_1-m_1-1}e_2^{r_2-1}T^{\alpha,\rightarrow}_{p_1-r_1-1-m_1,r_2-1}(r_1,r_2,s_1,s_2)\\
&=\Phi^{\alpha}(r_1,r_2)\Big\{\sum_{k_1=1}^{p_1-r_1-1}\frac{1}{\varphi_1^{-\alpha}(p_1-r_1,r_2)}T^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2)\\
&+\sum_{k_1=1}^{p_1-r_1-1}\frac{1}{\varphi_1^{-\alpha}(k_1,p_1-r_1,r_2)}\sum_{i_1=1}^{r_1-1}\frac{1}{\varphi_1^{\alpha}(i_1,r_1,r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2),
\end{aligned}
$$

and

$$
\begin{aligned}
&2p_1p_2\sum_{m_1=1}^{p_1-r_1-1}\sum_{m_2=1}^{p_2-r_2-1}\gamma^{\alpha}_{m_1}(r_1,r_2)\delta^{\alpha}_{m_2}(r_1,r_2)e_1^{p_1-1-m_1}e_2^{p_2-1-m_2}\\
&\qquad\qquad\times R^{\alpha,\rightarrow}_{p_1-r_1-1-m_1,p_2-r_2-1-m_2}(r_1,r_2,s_1,s_2)\\
&=\Phi^{\alpha}(r_1,r_2)\sum_{k_1=1}^{p_1-r_1-1}\frac{1}{\varphi_1^{-\alpha}(p_1-r_1,r_2)}\sum_{k_2=1}^{p_2-r_2-1}\frac{1}{\varphi_2^{-\alpha}(r_1,p_2-r_2)}B^{\alpha,\downarrow}_{0,0}(r_1,r_2,s_1,s_2).
\end{aligned}
$$

We finally have

$$
\begin{aligned}
&\Big(T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\Big)^2\\
&=T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)+\Psi_1^{\alpha}(r_1,r_2)T^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&+\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)+\Psi_1^{\alpha}(r_1,r_2)\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&-\Psi_1^{\alpha}(r_1,r_2)T^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)-\Psi_1^{\alpha}(r_1,r_2)\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&-\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)-\Psi_1^{\alpha}(r_1,r_2)\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&+\Psi_1^{\alpha}(r_1,r_2)\Psi_2^{\alpha}(r_1,r_2)B^{\alpha,\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)\\
&=T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2),
\end{aligned}
$$

which shows that $T^{\alpha}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)$ is idempotent. $\square$

5.2. **Block decomposition.** We can see that the elements

$$C_1=-q_1^{p_2}K^{-p_2}-q_1^{-p_2}K^{p_2}-(q_1^{p_2}-q_1^{-p_2})^2e_1f_1, \tag{5.2.1}$$

$$C_2=-q_2^{p_1}K^{-p_1}-q_2^{-p_1}K^{p_1}-(q_2^{p_1}-q_2^{-p_1})^2e_2f_2 \tag{5.2.2}$$

are central elements [5]. The central element C_1 acts as a scalar $\alpha^{p_2}(-1)^{r_2}(q_1^{p_2r_1}+q_1^{-p_2r_1})$ on the simple module $X^{\alpha}_{r_1,r_2}$ and the central element C_2 acts as a scalar $\alpha^{p_1(-1)^{r_1}}(q_2^{p_1r_2}+q_2^{-p_1r_2})$ on the simple module $X^{\alpha}_{r_1,r_2}$. We set $\beta_1(r_1,r_2)=\alpha^{p_2}(-1)^{r_2}(q_1^{p_2r_1}+q_1^{-p_2r_1})$ and $\beta_2^{\alpha}(r_1,r_2)=\alpha^{p_1}(-1)^{r_1}(q_2^{p_1r_2}+q_2^{-p_1r_2})$.

By direct calculations, we obtain the following.

Lemma 5.5. *We have*

$$
\begin{aligned}
&(C_1-\beta_1^{\alpha}(r_1,p_2))^2B^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,p_2,s_1,s_2)=0,\\
&(C_2-\beta_2^{\alpha}(r_1,p_2))B^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,p_2,s_1,s_2)=0,\\
&(C_1-\beta_1^{\alpha}(p_1,r_2))B^{\alpha,\uparrow}_{s_1-1,s_2-1}(p_1,r_2,s_1,s_2)=0,\\
&(C_2-\beta_2^{\alpha}(p_1,r_2))^2B^{\alpha,\uparrow}_{s_1-1,s_2-1}(p_1,r_2,s_1,s_2)=0,\\
&(C_1-\beta_1^{\alpha}(r_1,r_2))^2T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)=0,\\
&(C_2-\beta_2^{\alpha}(r_1,r_2))^2T^{\alpha,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2)=0.
\end{aligned}
$$

Set

$$
\begin{aligned}
&Q(p_1,p_2)=\{v\in\mathfrak{g}_{p_1,p_2}|(C_1-2)v=0,\ (C_2-2)v=0\},\\
&Q(0,p_2)=\{v\in\mathfrak{g}_{p_1,p_2}|(C_1-(-1)^{p_2}2)v=0,\ (C_2-(-1)^{p_1}2)v=0\},\\
&Q(r_1,p_2)=\left\{v\in\mathfrak{g}_{p_1,p_2}|(C_1-\beta_1^{+}(r_1,p_2))^2v=0,\ (C_2-(-1)^{p_1+r_1}2)v=0\right\},\\
&Q(p_1,r_2)=\left\{v\in\mathfrak{g}_{p_1,p_2}|(C_1-(-1)^{p_2+r_2}2)v=0,\ (C_2-\beta_2^{+}(p_1,r_2))^2v=0\right\},\\
&Q(r_1,r_2)=\left\{v\in\mathfrak{g}_{p_1,p_2}|(C_1-\beta_1^{+}(r_1,r_2))^2v=0,\ (C_2-\beta_2^{+}(r_1,r_2))^2v=0\right\}
\end{aligned}
$$

Since C_1 and C_2 are the central elements of $\mathfrak{g}_{p_1,p_2}$, we can see that the space $Q(r_1,r_2)$ is two-sided ideal of $\mathfrak{g}_{p_1,p_2}$. For $1\le r_1\le p_1-1$, we have $\beta_1^{+}(r_1,p_2)=\beta_1^{-}(p_1-r_1,p_2)$ and $\beta_2^{-}(p_1-r_1,p_2)=(-1)^{p_1}(-1)^{p_1-r_1}(-1)^{p_1}2=(-1)^{p_1+r_1}2$. Similarly we have $\beta_2^{+}(p_1,r_2)=\beta_2^{-}(p_1,p_2-r_2)$ and $\beta_1^{-}(p_1,p_2-r_2)=(-1)^{p}_{2}(-1)^{p_2-r_2}(-1)^{p_2}2=(-1)^{p_2+r_2}2$ for $1\le r_2\le p_2-1$. We also have

$$
\begin{aligned}
&\beta_1^{+}(r_1,r_2)=\beta_1^{-}(p_1-r_1,r_2)=\beta_1^{-}(r_1,p_2-r_1)=\beta_1^{+}(p_1-r_1,p_2-r_2),\\
&\beta_2^{+}(r_1,r_2)=\beta_2^{-}(p_1-r_1,r_2)=\beta_2^{-}(r_1,p_2-r_1)=\beta_2^{+}(p_1-r_1,p_2-r_2)
\end{aligned}
$$

for $1\le r_1\le p_1-1$ and $1\le r_2\le p_2-1$. By Lemma 5.5 we have injective maps

$$
\begin{aligned}
&X^{+}_{p_1,p_2}\hookrightarrow Q(p_1,p_2),\\
&X^{-}_{p_1,p_2}\hookrightarrow Q(0,p_2),\\
&P^{+}_{r_1,p_2},P^{-}_{p_1-r_1,p_2}\hookrightarrow Q(r_1,p_1),\\
&P^{+}_{p_1,r_2},P^{-}_{p_1,p_2-r_2}\hookrightarrow Q(p_1,r_2),\\
&P^{+}_{r_1,r_2},P^{-}_{r_1,p_2-r_2},P^{-}_{p_1-r_1,r_2},P^{+}_{p_1-r_1,p_2-r_2}\hookrightarrow Q(r_1,r_2).
\end{aligned}
$$

Note that $(\beta_1^{+}(r_1,p_2),(-1)^{p_1+r_1}2)=(\beta_1^{+}(r_1',p_2),(-1)^{p_1+r_1'}2)$ if and only if $r_1=r_1'$ for $1\le r_1,r_1'\le p_1-1$. We also have $(\beta_1^{+}(r_1,r_2),\beta_2^{+}(r_1,r_2))=(\beta_1^{+}(r_1',r_2'),\beta_2^{+}(r_1',r_2'))$ if and only if $(r_1',r_2')=(p_1-r_1,p_2-r_2),(r_1,r_2)$ for $1\le r_1,r_1'\le p_1-1$ and $1\le r_2,r_2'\le p_2-1$. Thus we have the decomposition of $\mathfrak{g}_{p_1,p_2}$ into two sided ideals.

Proposition 5.6 ([5])**.** *We have a decomposition of $\mathfrak{g}_{p_1,p_2}$ into two-sided ideals:*

$$
\mathfrak{g}_{p_1,p_2}=\bigoplus_{(r_1,r_2)\in I}Q(r_1,r_2)\oplus\bigoplus_{r_1=1}^{p_1-1}Q(r_1,p_2)\oplus\bigoplus_{r_2=1}^{p_2-1}Q(p_1,r_2)\oplus Q(p_1,p_2)\oplus Q(0,p_2)
$$

where

$$
I=\{(r_1,r_2)|1\le r_1\le p_1-1,1\le r_2\le p_2-1,p_2r_1+p_1r_2\le p_1p_2\}.
$$

Corollary 5.7. (1) *The elements* $B^{+,\downarrow}_{s_1-1,s_2-1}(p_1,p_2,s_1,s_2)$ *for* $1 \leq s_1 \leq p_1-1$ *and* $1 \leq s_2 \leq p_2-1$ *are mutually orthogonal primitive idempotents of* $Q(p_1,p_2)$.

(2) *The elements* $B^{-,\downarrow}_{s_1-1,s_2-1}(p_1,p_2,s_1,s_2)$, *for* $1 \leq s_1 \leq p_1-1$ *and* $1 \leq s_2 \leq p_2-1$, *are mutually orthogonal primitive idempotents of* $Q(0,p_2)$.

(3) *The elements*

$$\begin{aligned}
&B^{+,\uparrow}_{s_1-1,s_2-1}(r_1,p_2,s_1,s_2),\ 1 \leq s_1 \leq r_1,\ 1 \leq s_2 \leq p_2,\\
&B^{-,\uparrow}_{t_1-1,s_2-2}(p_1-r_1,p_2,t_1,s_2),\ 1 \leq t_1 \leq p_1-r_1,\ 1 \leq s_2 \leq p_2,
\end{aligned}$$

are mutually orthogonal primitive idempotents of $Q(r_1,p_2)$.

(4) *The elements*

$$\begin{aligned}
&B^{+,\uparrow}_{s_1-1,s_2-1}(p_1,r_2,s_1,s_2),\ 1 \leq s_1 \leq p_1,\ 1 \leq s_2 \leq r_2,\\
&B^{-,\uparrow}_{s_1-1,t_2-2}(p_1,p_2-r_2,s_1,t_2),\ 1 \leq s_1 \leq p_1,\ 1 \leq t_2 \leq p_2-r_2,
\end{aligned}$$

are mutually orthogonal primitive idempotents of $Q(p_1,r_2)$.

(5) *The elements*

$$\begin{aligned}
&T^{+,\uparrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2),\ 1 \leq s_1 \leq r_1,\ 1 \leq s_2 \leq r_2,\\
&T^{-,\uparrow}_{t_1-1,s_2-1}(p_1-r_1,r_2,t_1,s_2),\ 1 \leq t_1 \leq p_1-r_1,\ 1 \leq s_2 \leq r_2,\\
&T^{-,\uparrow}_{s_1-1,t_2-1}(r_1,p_2-r_2,s_1,t_2),\ 1 \leq s_1 \leq r_1,\ 1 \leq t_2 \leq p_2-r_2,\\
&T^{+,\uparrow}_{t_1-1,t_2-1}(p_1-r_1,p_2-r_2,t_1,t_2),\ 1 \leq t_1 \leq p_1-r_1,\ 1 \leq t_2 \leq p_2-r_2,
\end{aligned}$$

are mutually orthogonal primitive idempotents of $Q(r_1,r_2)$.

Proof. The corollary follows from Lemma 5.1, Proposition 5.2, Proposition 5.3, Proposition 5.4 and Lemma 5.5. ☐

Theorem 5.8. *The following equality holds*:

$$\begin{aligned}
&\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}\\
&=\bigoplus_{s_1=1}^{p_1}\bigoplus_{s_2=1}^{p_2} X^{+}_{p_1,p_2}(s_1,s_2) \oplus \bigoplus_{s_1=1}^{p_1}\bigoplus_{s_2=1}^{p_2} X^{-}_{p_1,p_2}(s_1,s_2)\\
&\oplus \bigoplus_{r_1=1}^{p_1-1}\bigoplus_{s_2=1}^{p_2}\left\{\bigoplus_{s_1=1}^{r_1} P^{\alpha}_{r_1,p_2}(s_1,s_2) \oplus \bigoplus_{t_1=1}^{p_1-r_1} P^{-}_{p_1-r_1,p_2}(t_1,s_2)\right\}\\
&\oplus \bigoplus_{r_2=1}^{p_2-1}\bigoplus_{s_1=1}^{p_1}\left\{\bigoplus_{s_2=1}^{r_2} P^{+}_{p_1,r_2}(s_1,s_2) \oplus \bigoplus_{t_2=1}^{p_2-r_2} P^{-}_{p_1,p_2-r_2}(s_1,t_2)\right\}\\
&\oplus \bigoplus_{(r_1,r_2)\in I}\left\{\bigoplus_{s_1=1}^{r_1}\bigoplus_{s_2=1}^{r_2} P^{+}_{r_1,r_2}(s_1,s_2) \oplus \bigoplus_{s_1=1}^{r_1}\bigoplus_{t_2=1}^{p_2-r_2} P^{-}_{r_1,p_2-r_2}(s_1,t_2)\right.\\
&\left.\oplus \bigoplus_{t_1=1}^{p_1-r_1}\bigoplus_{s_2=1}^{r_2} P^{-}_{p_1-r_1,r_2}(t_1,s_2) \oplus \bigoplus_{t_1=1}^{p_1-r_1}\bigoplus_{t_2=1}^{p_2-r_2} P^{+}_{p_1-r_1,p_2-r_2}(t_1,t_2)\right\}
\end{aligned}$$

Proof. By Corollary 5.7, the right hand side is contained in $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$. The dimension of the right hand side is

$$2p_1^2p_2^2 + 2p_1^2p_2^2(p_1-1) + 2p_1^2p_2^2(p_2-1) + 2p_1^2p_2^2(p_1-1)(p_2-1) = 2p_1^3p_2^3.$$

Since $\dim \mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}} = 2p_1^3p_2^3$, we have the equalitiy. □

6. MATRIX REALIZATION

Let A be a finite-dimensional associative algebra and $\{P_i \mid 1 \le i \le n\}$ the complete set of non-isomorphic indecomposable projective modules of A. Then the map defined by

$$A \to \bigoplus_{i=1}^{n} \mathrm{End}(P_i),\ \ a \mapsto (\rho_i(a))$$

is an algebra monomorphism where $\rho : A \to \mathrm{End}(P_i)$ is representation.

In this section, we give a matrix realization of each subalgebra Q_{r_1,r_2} of $\mathfrak{g}_{\boldsymbol{p_1},\boldsymbol{p_2}}$ by using the above fact.

6.1. **matrix realization of $Q(p_1,p_2)$ and $Q(0,p_2)$.** By Corollary 5.7 and Theorem 5.8, we have

$$Q(p_1,p_2) = \bigoplus_{s_1=1}^{p_1}\bigoplus_{s_2=1}^{p_2} X^{+}_{p_1,p_2}(s_1,s_2),$$
$$Q(0,p_2) = \bigoplus_{s_1=1}^{p_1}\bigoplus_{s_2=1}^{p_2} X^{-}_{p_1,p_2}(s_1,s_2).$$

By Lemma 5.1, the action of the basis of $B^{\alpha}_{n_1,n_2}(p_1,p_2,s_1,s_2)$ on the simple module $X^{\alpha}_{p_1,p_2}$ is given by

$$\begin{aligned} &B^{\alpha}_{m_1,m_2}(p_1,p_2,s_1,s_2)\mathsf{b}^{\alpha}_{n_1,n_2}(p_1,p_2) \\ &\qquad = \begin{cases} \mathsf{b}^{\alpha}_{m_1,m_2}(p_1,p_2), & (n_1,n_2)=(s_1-1,s_2-1), \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \tag{6.1.1}$$

Theorem 6.1. *The subalgebras $Q(p_1,p_2)$ and $Q(0,p_2)$ are isomorphic to the matrix algebra $M_{p_1p_2}(\mathbb{C})$. The isomorphism is given by*

$$B^{\alpha}_{m_1-1,m_2-1}(p_1,p_2,s_1,s_2) \mapsto E(m_1+p_1(m_2-1), s_1+p_1(s_2-1)),$$

for $1 \le m_i, s_i \le p_i$ where $E(a,b)$ is a matrix unit of $M_{p_1p_2}(\mathbb{C})$.

6.2. **matrix realization of $Q(r_1,p_2)$ and $Q(p_1,r_2)$.** By Corollary 5.7 and Theorem 5.8, we have

$$Q(r_1,p_2) = \bigoplus_{s_1=1}^{r_1}\bigoplus_{s_2=1}^{p_2} P^{+}_{r_1,p_2}(s_1,s_2) \oplus \bigoplus_{t_1=1}^{p_1-r_1}\bigoplus_{s_2=1}^{p_2} P^{-}_{p_1-r_1,p_2}(t_1,s_2),$$
$$Q(r_1,p_2) = \bigoplus_{s_1=1}^{p_1}\bigoplus_{s_2=1}^{r_2} P^{+}_{p_1,r_2}(s_1,s_2) \oplus \bigoplus_{s_1=1}^{p_1}\bigoplus_{t_2=1}^{p_2-r_2} P^{-}_{p_1,p_2-r_2}(s_1,t_2).$$

The action of $Q(r_1,p_2)$ on $P^{+}_{r_1,p_2}$ is give in the following table:

$Q(r_1,p_2)\backslash P^{+}_{r_1,p_2}$	$\mathsf{b}^{+,\uparrow}_{s_1-1,s_2-1}$	$\mathsf{b}^{+,\leftarrow}_{t_1-1,s_2-1}$	$\mathsf{b}^{+,\rightarrow}_{t_1-1,s_2-1}$	$\mathsf{b}^{+,\downarrow}_{s_1-1,s_2-1}$
$B^{+,\uparrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)$	$\mathsf{b}^{+,\uparrow}_{n_1,n_2}$	0	0	$\mathsf{b}^{+,\downarrow}_{n_1,n_2}$
$B^{+,\leftarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)$	$\mathsf{b}^{+,\leftarrow}_{k_1,n_2}$	0	0	0
$B^{+,\rightarrow}_{k_1,n_2}(r_1,p_2,s_1,s_2)$	$\mathsf{b}^{+,\rightarrow}_{k_1,b_2}$	0	0	0
$B^{+,\downarrow}_{n_1,n_2}(r_1,p_2,s_1,s_2)$	$\mathsf{b}^{+,\downarrow}_{n_1,n_2}$	0	0	0
$B^{-,\uparrow}_{k_1,n_2}(p_1-r_1,p_2,t_1,s_2)$	0	$\mathsf{b}^{+,\leftarrow}_{k_1,n_2}$	$\mathsf{b}^{+,\rightarrow}_{k_1,2}$	0
$B^{-,\leftarrow}_{n_1,n_2}(p_1-r_1,p_2,t_1,s_2)$	0	0	$\mathsf{b}^{+,\downarrow}_{n_1,n_2}$	0
$B^{-,\rightarrow}_{n_1,n_2}(p_1-r_1,p_2,t_1,s_2)$	0	$\mathsf{b}^{+,\downarrow}_{n_1,n_2}$	0	0
$B^{-,\downarrow}_{k_1,n_2}(p_1-s_1,p_2,t_1,s_2)$	0	0	0	0

Remark that the actions which do not appear in the above table are all zero. Substituting $+, r_1$ with $-, p_1-r_1$ in above table, we have the actions of the basis of $Q(r_1,p_2)$ on $P^{-}_{p_1-r_1,p_2}$.

Let us expand $v \in Q(r_1,p_2)$ as follows:

$$
\begin{aligned}
v = &\sum_{s_1=1}^{r_1}\sum_{s_2=1}^{p_2}\sum_{n_2=1}^{p_2}\Big\{\sum_{\bullet\in\{\uparrow,\downarrow\}}\sum_{n_1=1}^{r_1} b^{+,\bullet}_{r_1,p_2}(n_1+(n_2-1)r_1, s_1+(s_2-1)r_1)(v)\\
&B^{+,\bullet}_{n_1-1,n_2-1}(r_1,p_2,s_1,s_2)\\
&+\sum_{\bullet\in\{\leftarrow,\rightarrow\}}\sum_{k_1=1}^{p_1-r_1} b^{+,\bullet}_{r_1,p_2}(k_1+(n_2-1)(p_1-r_1), s_1+(s_2-1)r_1)(v)\\
&B^{+,\bullet}_{k_1-1,n_2-1}(r_1,p_2,s_1,s_2)\Big\}\\
&+\sum_{t_1=1}^{p_1-r_1}\sum_{s_2=1}^{p_2}\sum_{n_2=1}^{p_2}\Big\{\sum_{\bullet\in\{\uparrow,\downarrow\}}\sum_{k_1=1}^{p_1-r_1} b^{-,\bullet}_{p_1-r_1,p_2}(k_1+(n_2-1)(p_1-r_1), s_1+(s_2-1)(p_1-r_1))(v)\\
&\times B^{+,\bullet}_{k_1-1,n_2-1}(p_1-r_1,p_2,t_1,s_2)\\
&+\sum_{\bullet\in\{\leftarrow,\rightarrow\}}\sum_{n_1=1}^{r_1} b^{-,\bullet}_{p_1-r_1,p_2}(n_1+(n_2-1)r_1, s_1+(s_2-1)r_1)(v)\\
&\times B^{;\bullet}_{n_1-1,n_2-1}(p_1-r_1,p_2,t_1,s_2)\Big\}.
\end{aligned}
$$

We define

$$
\begin{aligned}
B^{+,\bullet}_{r_1,p_2}(v) &= [b^{+,\bullet}_{r_1,p_2}(m_1,m_2)(v)] \in M_{r_1p_2}(\mathbb{C}),\\
B^{+,\circ}_{r_1,p_2}(v) &= [b^{+,\bullet}_{r_1,p_2}(m_1,m_2)(v)] \in M_{(p_1-r_1)p_2,r_1p_2}(\mathbb{C}),\\
B^{-,\bullet}_{p_1-r_1,p_2} &= [b^{-,\bullet}_{p_1-r_1,p_2}(m_1,m_2)(v)] \in M_{(p_1-r_1)p_2}(\mathbb{C}),\\
B^{-,\circ}_{p_1-r_1,p_2} &= [b^{-,\circ}_{p_1-r_1,p_2}(m_1,m_2)(v)] \in M_{r_1p_2,(p_1-r_1)p_2}(\mathbb{C})
\end{aligned}
$$

where $\bullet \in \{\uparrow,\downarrow\}$ and $\circ \in \{\leftarrow,\rightarrow\}$. Then, by the action of $Q(r_1,p_2)$ on the projective modules $P^{+}_{r_1,p_2}$ and $P^{-}_{p_1-r_1,p_2}$, we have the following result.

Theorem 6.2. *For* $1 \leq r_1 \leq p_1 - 1$, $Q(r_1, p_2)$ *is isomorphic to the subalgebra of* $M_{2p_1p_2}(\mathbb{C}) \oplus M_{2p_1p_2}(\mathbb{C})$ *with the shape:*

$$\left(\begin{bmatrix} B^{+,\uparrow}_{r_1,p_2} & 0 & 0 & 0 \\ B^{+,\leftarrow}_{r_1,p_2} & B^{-,\uparrow}_{p_1-r_1,p_2} & 0 & 0 \\ B^{+,\rightarrow}_{r_1,p_2} & 0 & B^{-,\uparrow}_{p_1-r_1,p_2} & 0 \\ B^{+,\downarrow}_{r_1,p_2} & B^{-,\rightarrow}_{p_1-r_1,p_2} & B^{-,\leftarrow}_{p_1-r_1,p_2} & B^{+,\uparrow}_{r_1,p_2} \end{bmatrix}, \begin{bmatrix} B^{-,\uparrow}_{p_1-r_1,p_2} & 0 & 0 & 0 \\ B^{-,\leftarrow}_{p_1-r_1,p_2} & B^{+,\uparrow}_{r_1,p_2} & 0 & 0 \\ B^{-,\rightarrow}_{p_1-r_1,p_2} & 0 & B^{+,\uparrow}_{r_1,p_2} & 0 \\ B^{-,\downarrow}_{p_1-r_1,p_2} & B^{+,\rightarrow}_{r_1,p_2} & B^{-,\leftarrow}_{r_1,p_2} & B^{-,\uparrow}_{p_1-r_1,p_2} \end{bmatrix} \right).$$

By using similar argument for the case $Q(r_1, p_2)$, we have the following:

Theorem 6.3. *For* $1 \leq r_1 \leq p_1 - 1$, *the algebra* $Q(r_1, p_2)$ *is isomorphic to the subalgebra of* $M_{2p_1p_2}(\mathbb{C}) \oplus M_{2p_1p_2}(\mathbb{C})$*:*

$$\left(\begin{bmatrix} B^{+,\uparrow}_{p_1,r_2} & 0 & 0 & 0 \\ B^{+,\leftarrow}_{p_1,r_2} & B^{-,\uparrow}_{p_1,p_2-r_2} & 0 & 0 \\ B^{+,\rightarrow}_{p_1,r_2} & 0 & B^{-,\uparrow}_{p_1,p_2-r_2} & 0 \\ B^{+,\downarrow}_{p_1,r_2} & B^{-,\rightarrow}_{p_1,p_2-r_2} & B^{-,\leftarrow}_{p_1,p_2-r_2} & B^{+,\uparrow}_{p_1,r_2} \end{bmatrix}, \begin{bmatrix} B^{-,\uparrow}_{p_1,p_2-r_2} & 0 & 0 & 0 \\ B^{-,\leftarrow}_{p_1,p_2-r_2} & B^{+,\uparrow}_{p_1,r_2} & 0 & 0 \\ B^{-,\rightarrow}_{p_1,p_2-r_2} & 0 & B^{+,\uparrow}_{p_1,r_2} & 0 \\ B^{-}_{p_1,p_2-r_2} & B^{+,\rightarrow}_{p_1,r_2} & B^{+,\leftarrow}_{p_1,r_2} & B^{-,\uparrow}_{p_1,p_2-r_2} \end{bmatrix} \right).$$

6.3. **matrix realization of $Q(r_1, r_2)$.** By Corollary 5.7 and Theorem 5.8, we have

$$\begin{aligned} Q(r_1, r_2) = & \bigoplus_{s_1=1}^{r_1} \bigoplus_{s_2=1}^{r_2} P^{+}_{r_1,r_2}(s_1, s_2) \oplus \bigoplus_{s_1=1}^{r_1} \bigoplus_{t_2=1}^{p_2-r_2} P^{-}_{r_1,p_2-r_2}(s_1, t_2) \\ & \oplus \bigoplus_{t_1=1}^{p_1-r_1} \bigoplus_{s_2=1}^{r_2} P^{-}_{p_1-r_1,r_2}(t_1, s_2) \oplus \bigoplus_{t_1=1}^{p_1-r_1} \bigoplus_{t_2=1}^{p_2-r_2} P^{+}_{p_1-r_1,p_2-r_2}(t_1, t_2). \end{aligned}$$

Since $Q(r_1, r_2)$ has four non-isomorphic indecomposable projective modules, there is an algebra monomorphism

$$Q(r_1, r_2) \to \mathrm{End}_{\mathbb{C}}(P^{+}_{r_1,r_2}) \oplus \mathrm{End}_{\mathbb{C}}(P^{-}_{r_1,p_2-r_2}) \oplus \mathrm{End}_{\mathbb{C}}(P^{-}_{p_1-r_1,r_2}) \oplus \mathrm{End}_{\mathbb{C}}(P^{+}_{p_1-r_1,p_2-r_2}).$$

By Lemma 5.1, the definition of the basis of $Q(r_1, r_2)$ and the structure of projective modules, we can determine the action of $Q(r_1, r_2)$ on the projective module as in Table 1 (the actions do not appear in Table 1 are zero).

Table 1. The actions of the basis of $Q(r_1,r_2)$ on $P^+_{r_1,r_2}$.

$P^+_{r_1,r_2}(s_1,s_2)\backslash P^+_{r_1,r_2}$	$\mathsf{t}^{+,\uparrow}_{s_1-1,s_2-1}$	$\mathsf{t}^{+,\downarrow}_{s_1-1,s_2-1}$	$\mathsf{b}^{+,\uparrow}_{s_1-1,s_2-1}$	$\mathsf{b}^{+,\downarrow}_{s_1-1,s_2-1}$
$T^{+,\uparrow}_{m_1,m_2}(r_1,r_2,s_1,s_2)$	$\mathsf{t}^{+,\uparrow}_{m_1,m_2}$	$\mathsf{t}^{+,\downarrow}_{m_1,m_2}$	$\mathsf{b}^{+,\uparrow}_{m_1,m_2}$	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$
$T^{+,\bullet}_{k_1,n_2}(r_1,r_2,s_1,s_2)$	$\mathsf{t}^{+,\bullet}_{k_1,n_2}$	0	$\mathsf{b}^{+,\bullet}_{k_1,m_2}$	0
$X^{+,\uparrow}_{m_1,k_2}(r_1,r_2,s_1,s_2)$	$\mathsf{x}^{+,\uparrow}_{m_1,k_2}$	$\mathsf{x}^{+,\downarrow}_{m_1,k_2}$	0	0
$X^{+,\bullet}_{m_1,k_2}(r_1,r_2,s_1,s_2)$	$\mathsf{x}^{+,\bullet}_{m_1,k_2}$	0	0	0
$B^{+,\uparrow}_{m_1,m_2}(r_1,r_2,s_1,s_2)$	$\mathsf{b}^{+,\uparrow}_{m_1,m_2}$	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0	0
$B^{+,\bullet}_{k_1,n_2}(r_1,r_2,s_1,s_2)$	$\mathsf{b}^{+,\bullet}_{k_1,m_2}$	0	0	0

$P^-_{p_1-r_1,r_2}(t_1,s_2)\backslash P^+_{r_1,r_2}$	$\mathsf{t}^{+,\leftarrow}_{t_1-1,s_2-1}$	$\mathsf{t}^{+,\rightarrow}_{t_1-1,s_2-1}$	$\mathsf{b}^{+,\leftarrow}_{t_1-1,s_2-1}$	$\mathsf{b}^{+,\rightarrow}_{t_1-1,s_2-1}$
$T^{-,\uparrow}_{k_1,m_2}(p_1-r_1,r_2,t_1,s_2)$	$\mathsf{t}^{+,\leftarrow}_{k_1,m_2}$	$\mathsf{t}^{+,\rightarrow}_{k_1,m_2}$	$\mathsf{b}^{+,\leftarrow}_{k_1,m_2}$	$\mathsf{b}^{+,\rightarrow}_{k_1,m_2}$
$T^{-,\leftarrow}_{m_1,m_2}(p_1-r_1,r_2,t_1,s_2)$	0	$\mathsf{t}^{+,\downarrow}_{m_1,m_2}$	0	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$
$T^{-,\rightarrow}_{m_1,m_2}(p_1-r_1,r_2,t_1,s_2)$	$\mathsf{t}^{+,\downarrow}_{m_1,m_2}$	0	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0
$X^{-,\uparrow}_{k_1,k_2}(p_1-r_1,r_2,t_1,s_2)$	$\mathsf{x}^{+,\leftarrow}_{k_1,k_2}$	$\mathsf{x}^{+,\rightarrow}_{k_1,k_2}$	0	0
$X^{-,\leftarrow}_{m_1,k_2}(p_1-r_1,r_2,t_1,s_2)$	0	$\mathsf{x}^{+,\downarrow}_{m_1,k_2}$	0	0
$X^{-,\rightarrow}_{m_1,k_2}(p_1-r_1,r_2,t_1,s_2)$	$\mathsf{x}^{+,\downarrow}_{m_1,k_2}$	0	0	0

$P^-_{r_1,p_2-r_2}(s_1,t_2)\backslash P^+_{r_1,r_2}$	$\mathsf{l}^{+,\uparrow}_{s_1-1,t_2-1}$	$\mathsf{l}^{+,\downarrow}_{s_1-1,t_2-1}$	$\mathsf{r}^{+,\uparrow}_{s_1-1,t_2-1}$	$\mathsf{r}^{+,\downarrow}_{s_1-1,t_2-1}$
$T^{-,\uparrow}_{m_1,k_2}(r_1,p_2-r_2,s_1,t_2)$	$\mathsf{l}^{+,\uparrow}_{m_1,k_2}$	$\mathsf{l}^{+,\downarrow}_{m_1,k_2}$	$\mathsf{r}^{+,\uparrow}_{m_1,k_2}$	$\mathsf{r}^{+,\downarrow}_{m_1,k_2}$
$T^{-,\bullet}_{k_1,k_2}(r_1,p_2-r_2,s_1,t_2)$	$\mathsf{l}^{+,\bullet}_{k_1,k_2}$	0	$\mathsf{r}^{+,\bullet}_{k_1,k_2}$	0
$L^{-,\uparrow}_{m_1,m_2}(r_1,p_2-r_2,s_1,t_2)$	0	0	$\mathsf{b}^{+,\uparrow}_{m_1,m_2}$	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$
$L^{-,\bullet}_{k_1,m_2}(r_1,p_2-r_2,s_1,t_2)$	0	0	$\mathsf{b}^{+,\bullet}_{k_1,m_2}$	0
$R^{-,\uparrow}_{m_1,m_2}(r_1,p_2-r_2,s_1,t_2)$	$\mathsf{b}^{+,\uparrow}_{m_1,m_2}$	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0	0
$R^{-,\bullet}_{k_1,m_2}(r_1,p_2-r_2,s_1,t_2)$	$\mathsf{b}^{+,\bullet}_{k_1,m_2}$	0	0	0

$P^+_{p_1-r_1,p_2-r_2}(t_1,t_2)\backslash P^+_{r_1,r_2}$	$\mathsf{l}^{+,\leftarrow}_{t_1-1,t_2-1}$	$\mathsf{l}^{+,\rightarrow}_{t_1-1,t_2-1}$	$\mathsf{r}^{+,\leftarrow}_{t_1-1,r_2-1}$	$\mathsf{r}^{+,\rightarrow}_{t_1-1,t_2-1}$
$T^{+,\uparrow}_{k_1,k_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	$\mathsf{l}^{+,\leftarrow}_{k_1,k_2}$	$\mathsf{l}^{+,\rightarrow}_{k_1,k_2}$	$\mathsf{r}^{+,\leftarrow}_{k_1,k_2}$	$\mathsf{r}^{+,\rightarrow}_{k_1,k_2}$
$T^{+,\leftarrow}_{m_1,k_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	0	$\mathsf{l}^{+,\downarrow}_{m_1,k_2}$	0	$\mathsf{r}^{+,\downarrow}_{m_1,k_2}$
$T^{+,\rightarrow}_{m_1,k_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	$\mathsf{l}^{+,\downarrow}_{m_1,k_2}$	0	$\mathsf{r}^{+,\downarrow}_{m_1,k_2}$	0
$L^{+,\uparrow}_{k_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	0	0	$\mathsf{b}^{+,\leftarrow}_{k_1,m_2}$	$\mathsf{b}^{+,\rightarrow}_{k_1,m_2}$
$L^{+,\leftarrow}_{m_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	0	0	0	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$
$L^{+,\rightarrow}_{m_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	0	0	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0
$R^{+,\uparrow}_{k_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	$\mathsf{b}^{+,\leftarrow}_{k_1,m_2}$	$\mathsf{b}^{+,\rightarrow}_{k_1,m_2}$	0	0
$R^{+,\leftarrow}_{m_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	0	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0	0
$R^{+,\rightarrow}_{m_1,m_2}(p_1-r_1,p_2-r_2,t_1,t_2)$	$\mathsf{b}^{+,\downarrow}_{m_1,m_2}$	0	0	0

Let $\rho^+_{r_1,r_2}$ be the representation of $Q(r_1,r_2)$ on $P^+_{r_1,r_2}$. Let us expand $v\in Q(r_1,r_2)$ as

$$v=v^+_{r_1,r_2}+v^-_{p_1-r_1,r_2}+v^-_{r_1,p_2-r_2}+v^+_{p_1-r_1,p_2-r_2}$$

where

$$
\begin{aligned}
&v^{+}_{r_1,r_2} = \sum_{s_1=1}^{r_1}\sum_{s_2=1}^{r_2} \\
&\Big\{ \sum_{\bullet\in\{\uparrow,\downarrow\}} \sum_{n_1=1}^{r_1}\sum_{n_2=1}^{r_2} \Big(t^{+,\bullet}_{r_1,r_2}(n_1+r_1(n_2-1), s_1+(s_2-1)r_2)(v)T^{+,\bullet}_{n_1-1,n_2-1}(r_1,r_2,s_1,s_2) \\
&+ b^{+,\bullet}_{r_1,r_2}(n_1+r_1(n_2-1), s_1+(s_2-1)r_1)(v)T^{+,\bullet}_{n_1-1,n_2-1}(r_1,r_2,s_1,s_2)\Big) \\
&+ \sum_{\bullet\in\{\leftarrow,\rightarrow\}} \sum_{k_1=1}^{p_1-r_1}\sum_{n_2=1}^{r_2} \\
&\Big(t^{+,\bullet}_{r_1,r_2}(k_1+(p_1-r_1)(n_2-1), s_1+(s_2-1)r_1)(v)T^{+,\bullet}_{k_1-1,n_2-1}(r_1,r_2,s_1,s_2) \\
&+ b^{+,\bullet}_{r_1,r_2}(k_1+(p_1-r_1)(n_2-1), s_1+(s_2-1)r_1)(v)B^{+,\bullet}_{k_1-1,n_2-1}(r_1,r_2,s_1,s_2)\Big) \\
&\sum_{\bullet\in\{\uparrow,\downarrow\}} \sum_{n_1=1}^{r_1}\sum_{k_2=1}^{p_2-r_2} \Big(r^{+,\bullet}_{r_1,r_2}(n_1+r_1(k_2-1), s_1+(s_2-1)r_1)(v)R^{+,\bullet}_{n_1-1,k_2-1}(r_1,r_2,s_1,s_2) \\
&+ b^{+,\bullet}_{r_1,r_2}(n_1+r_1(k_2-1), s_1+(s_2-1)r_1)(v)L^{+,\bullet}_{n_1-1,k_2-1}(r_1,r_2,s_1,s_2)\Big) \\
&+ \sum_{\bullet\in\{\leftarrow,\rightarrow\}} \sum_{k_1=1}^{p_1-r_1}\sum_{k_2=1}^{p_2-r_2} \\
&\Big(r^{+,\bullet}_{r_1,r_2}(k_1+(p_1-r_1)(k_2-1), s_1+(s_2-1)r_1)(v)R^{+,\bullet}_{k_1-1,k_2-1}(r_1,r_2,s_1,s_2) \\
&+ b^{+,\bullet}_{r_1,r_2}(k_1+(p_1-r_1)(k_2-1), s_1+(s_2-1)r_1)(v)L^{+,\bullet}_{k_1-1,k_2-1}(r_1,r_2,s_1,s_2)\Big)\Big\}
\end{aligned}
$$

and set

$$
\begin{aligned}
&T^{+,\bullet}_{r_1,r_2}(v) = [t^{+,\bullet}_{r_1,r_2}(m_1,m_2)(v)], B^{+,\bullet}_{r_1,r_2}(v) = [b^{+,\bullet}_{r_1,r_2}(m_1,m_2)(v)] \in M_{r_1r_2}(\mathbb{C}), \\
&T^{+,\circ}_{r_1,r_2}(v) = [t^{+,\circ}_{r_1,r_2}(m_1,m_2)(v)], B^{+,\circ}_{r_1,r_2}(v) = [b^{+,\circ}_{r_1,r_2}(m_1,m_2)(v)] \in M_{(p_1-r_1)r_2,r_1r_2}(\mathbb{C}), \\
&R^{+,\bullet}_{r_1,r_2}(v) = [r^{+,\bullet}_{r_1,r_2}(m_1,m_2)(v)], L^{+,\bullet}_{r_1,r_2} = [l^{+,\bullet}_{r_1,r_2}(m_1,m_2)(v)] \in M_{r_1(p_2-r_2),r_1r_2}(\mathbb{C}), \\
&R^{+,\circ}_{r_1,r_2}(v) = [r^{+,\circ}_{r_1,r_2}(m_1,m_2)(v)], L^{+,\circ}_{r_1,r_2}(v) = [l^{+,\circ}_{r_1,r_2}(m_1,m_2)(v)] \in M_{(p_1-r_1)(p_2-r_2),r_1r_2}(\mathbb{C}),
\end{aligned}
$$

for $\bullet \in \{\uparrow,\downarrow\}$ and $\circ \in \{\leftarrow,\rightarrow\}$. Substituting $(+,r_1,r_2)$ with $(-,p_1-r_1,r_2)$, $(-,r_1,p_2-r_2)$ and $(+,p_1-r_1,p_2-r_2)$, we can define $v^{-}_{p_1-r_1,r_2}$, $v^{-}_{r_1,p_2-r_2}$ and $v^{+}_{p_1-r_1,p_2-r_2}$ and corresponding matrices. By Table 1, the linear map $\rho^{+}_{r_1,r_2}(v)$ for $v \in Q(r_1,r_2)$ can be expressed as follows

$$
\begin{bmatrix}
T^{+}_{r_1,r_2}(v) & 0 & 0 & 0 \\
L^{+}_{r_1,r_2}(v) & T^{-}_{r_1,p_2-r_2}(v) & 0 & 0 \\
R^{+}_{r_1,r_2}(v) & 0 & T^{-}_{r_1,p_2-r_2}(v) & 0 \\
B^{+}_{r_1,r_2}(v) & R^{-}_{r_1,p_2-r_2}(v) & L^{-}_{r_1,p_2-r_2}(v) & T^{+}_{r_1,r_2}(v)
\end{bmatrix}
$$

with

$$X^{+}_{r_1,r_2}(v) = \begin{bmatrix} X^{+,\uparrow}_{r_1,r_2}(v) & 0 & 0 & 0 \\ X^{+,\leftarrow}_{r_1,r_2}(v) & X^{-,\uparrow}_{p_1-r_1,r_2}(v) & 0 & 0 \\ X^{+,\rightarrow}_{r_1,r_2}(v) & 0 & X^{-,\uparrow}_{p_1-r_1,r_2}(v) & 0 \\ X^{+,\downarrow}_{r_1,r_2}(v) & X^{-,\rightarrow}_{p_1-r_1,r_2}(v) & X^{-,\leftarrow}_{p_1-r_1,r_2}(v) & X^{+,\uparrow}_{r_1,r_2}(v) \end{bmatrix},$$

where $X \in \{T, L, R, B\}$.

Then we have the following result.

Theorem 6.4. *The algebra of $Q(r_1, r_2)$ with $(r_1, r_2) \in I$ is isomorphic to the subalgebra of $M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C}) \oplus M_{4p_1p_2}(\mathbb{C})$ given by*

$$\left(\begin{bmatrix} T^{+}_{r_1,r_2} & 0 & 0 & 0 \\ L^{+}_{r_1,r_2} & T^{-}_{r_1,p_2-r_2} & 0 & 0 \\ R^{+}_{r_1,r_2} & 0 & T^{-}_{r_1,p_2-r_2} & 0 \\ B^{+}_{r_1,r_2} & R^{-}_{r_1,p_2-r_2} & L^{-}_{r_1,p_2-r_2} & T^{+}_{r_1,r_2} \end{bmatrix}, \begin{bmatrix} T^{-}_{p_1-r_1,r_2} & 0 & 0 & 0 \\ L^{-}_{p_1-r_1,r_2} & T^{+}_{p_1-r_1,p_2-r_2} & 0 & 0 \\ R^{-}_{p_1-r_1,r_2} & 0 & T^{+}_{p_1-r_1,p_2-r_2} & 0 \\ B^{-}_{p_1-r_1,r_2} & R^{+}_{p_1-r_1,p_2-r_2} & L^{+}_{p_1-r_1,p_2-r_2} & T^{-}_{p_1-r_1,r_2} \end{bmatrix}, \right.$$

$$\left. \begin{bmatrix} T^{-}_{r_1,p_2-r_2} & 0 & 0 & 0 \\ L^{-}_{r_1,p_2-r_2} & T^{+}_{r_1,r_2} & 0 & 0 \\ R^{-}_{r_1,p_2-r_2} & 0 & T^{+}_{r_1,r_2} & 0 \\ B^{-}_{r_1,p_2-r_2} & R^{+}_{r_1,r_2} & L^{+}_{r_1,r_2} & T_{r_1,p_2-r_2} \end{bmatrix}, \begin{bmatrix} T^{+}_{p_1-r_1,p_2-r_2} & 0 & 0 & 0 \\ L^{+}_{p_1-r_1,p_2-r_2} & T^{-}_{p_1-r_1,r_2} & 0 & 0 \\ R^{+}_{p_1-r_1,p_2-r_2} & 0 & T^{-}_{p_1-r_1,r_2} & 0 \\ B^{+}_{p_1-r_1,p_2-r_2} & R^{-}_{p_1-r_1,r_2} & L^{-}_{p_1-r_1,r_2} & T^{+}_{p_1-r_1,p_2-r_2} \end{bmatrix} \right)$$

where

$$X^{\alpha}_{r_1,r_2} = \begin{bmatrix} X^{\alpha,\uparrow}_{r_1,r_2} & 0 & 0 & 0 \\ X^{\alpha,\leftarrow}_{r_1,r_2} & X^{-\alpha,\uparrow}_{p_1-r_1,r_2} & 0 & 0 \\ X^{\alpha,\rightarrow}_{r_1,r_2} & 0 & X^{-\alpha,\uparrow}_{p_1-r_1,r_2} & 0 \\ X^{\alpha,\downarrow}_{r_1,r_2} & X^{-\alpha,\rightarrow}_{p_1-r_1,r_2} & X^{-\alpha,\leftarrow}_{p_1-r_1,r_2} & X^{\alpha,\uparrow}_{r_1,r_2} \end{bmatrix}$$

and $X = T, R, L, B$.

7. SYMMETRIC LINEAR FUNCTIONS ON $\mathfrak{g}_{p_1,p_2}$

7.1. **The center.** The structure of the center of $\mathfrak{g}_{p_1,p_2}$ is described in [5]. Recall that $\mathfrak{g}_{p_1,p_2}$ decomposes into subalgebras as

$$\mathfrak{g}_{p_1,p_2} = \bigoplus_{(r_1,r_2)\in I} Q(r_1,r_2) \oplus \bigoplus_{r_1=1}^{p_1-1} Q(r_1,p_2) \oplus \bigoplus_{r_2=1}^{p_2-1} Q(p_1,r_2) \oplus Q(p_1,p_2) \oplus Q(0,p_2).$$

Proposition 7.1 ([5])**.**

$$\dim Z(Q(r_1,r_2)) = \begin{cases} 9, & (r_1,r_2) \in I, \\ 3, & 1 \le r_1 \le p_1 - 1, r_2 = p_2, \\ 3, & r_1 = p_1, 1 \le r_2 \le p_2 - 1, \\ 1, & (r_1,r_2) = (p_1,p_2). \end{cases}$$

By Proposition 2.2 we can see that the space of symmetric linear functions on $\mathfrak{g}_{p_1,p_2}$ is $\frac{1}{2}(3p_1 - 1)(3p_2 - 1)$-dimensional.

The action of the center on the projective modules is also determined in [5]. For each $Q(r_1, r_2)$, there is a central idempotent $e(r_1, r_2)$ which acts as an identity on $Q(r_1, r_2)$ and as zero on $Q(s_1, s_2)$ for $(r_1, r_2) \neq (s_1, s_2)$. For $Q(r_1, r_2)$ with $(r_1, r_2) \in I$, there are eight central elements as follows;

$$\begin{aligned}
&v^{\nearrow}(r_1,r_2),\ v^{\nearrow}(r_1,r_2)\mathsf{t}^{\bullet}_{n_1,n_2} = \mathsf{b}^{\bullet}_{n_1,n_2},\ \bullet \in \{\uparrow,\downarrow,\leftarrow,\rightarrow\} \text{ in } P^{+}_{r_1,r_2} \text{ and } P^{-}_{p_1-r_1,r_2},\\
&v^{\nwarrow}(r_1,r_2),\ v^{\nwarrow}(r_1,r_2)\mathsf{x}^{\uparrow}_{n_1,n_2} = \mathsf{x}^{\downarrow}_{n_1,n_2},\ \mathsf{x} \in \{\mathsf{t},\mathsf{b},\mathsf{l},\mathsf{r}\} \text{ in } P^{+}_{r_1,r_2} \text{ and } P^{-}_{r_1,p_2-r_2},\\
&v^{\swarrow}(r_1,r_2),\ v^{\swarrow}(r_1,r_2)\mathsf{t}^{\bullet}_{n_1,n_2} = \mathsf{b}^{\bullet}_{n_1,n_2},\ \bullet \in \{\uparrow,\downarrow,\leftarrow,\rightarrow\} \text{ in } P^{-}_{r_1,p_2-r_2} \text{ and } P^{+}_{p_1-r_1,p_2-r_2},\\
&v^{\searrow}(r_1,r_2),\ v^{\searrow}(r_1,r_2)\mathsf{x}^{\uparrow}_{n_1,n_2} = \mathsf{x}^{\downarrow}_{n_1,n_2},\ \mathsf{x} \in \{\mathsf{t},\mathsf{b},\mathsf{l},\mathsf{r}\} \text{ in } P^{-}_{p_1-r_1,r_2} \text{ and } P^{+}_{p_1-r_1,p_2-r_2},\\
&w^{\uparrow}(r_1,r_2),\ w^{\uparrow}(r_1,r_2)\mathsf{t}^{\uparrow}_{n_1,n_2} = \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{+}_{r_1,r_2},\\
&w^{\rightarrow}(r_1,r_2),\ w^{\rightarrow}(r_1,r_2)\mathsf{t}^{\uparrow}_{n_1,n_2} = \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{-}_{p_1-r_1,r_2},\\
&w^{\leftarrow}(r_1,r_2),\ w^{\leftarrow}(r_1,r_2)\mathsf{t}^{\uparrow}_{n_1,n_2} = \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{-}_{r_1,p_2-r_2},\\
&w^{\downarrow}(r_1,r_2),\ w^{\downarrow}(r_1,r_2)\mathsf{t}^{\uparrow}_{n_1,n_2} = \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{+}_{p_1-r_1,p_2-r_2}.
\end{aligned}$$

There are two central elements $w^{\pm}(r_1, p_2)$ in $Q(r_1, p_2)$ and we have

$$\begin{aligned}
w^{+}(r_1,p_2)\mathsf{b}^{\uparrow}_{n_1,n_2} &= \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{+}_{r_1,p_2},\\
w^{-}(r_1,p_2)\mathsf{b}^{\uparrow}_{n_1,n_2} &= \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{-}_{p_1-r_1,p_2}.
\end{aligned}$$

Similarly we can determine the actions of the central elements $w^{\pm}(p_1, r_2)$ in $Q(p_1, r_2)$:

$$\begin{aligned}
w^{+}(p_1,r_2)\mathsf{b}^{\uparrow}_{n_1,n_2} &= \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{+}_{p_1,r_2},\\
w^{-}(p_1,r_2)\mathsf{b}^{\uparrow}_{n_1,n_2} &= \mathsf{b}^{\downarrow}_{n_1,n_2} \text{ in } P^{-}_{p_1,p_2-r_2}.
\end{aligned}$$

7.2. **Symmetric linear functions.** By Theorem 6.1, the subalgebras $Q(p_1, p_2)$ and $Q(0, p_2)$ are isomorphic to the matrix algebra $M_{p_1p_2}(\mathbb{C})$. Thus we can define the linear functions τ_{p_1,p_2} on $Q(p_1, p_2)$ and τ_{0,p_2} on $Q(0, p_2)$ as the trace on $M_{p_1p_2}(\mathbb{C})$. It is clear that the linear functions τ_{p_1,p_2} and τ_{0,p_2} are symmetric.

By Theorem 6.2, we define the linear functions on $Q(r_1, p_2)$ by

$$\begin{aligned}
&\tau^{+}_{r_1,p_2}(v) = \operatorname{tr} B^{+,\uparrow}_{r_1,p_2}(v),\ \tau^{-}_{r_1,p_2}(v) = \operatorname{tr} B^{-,\uparrow}_{p_1-r_1,p_2}(v),\\
&\chi_{r_1,p_2}(v) = \operatorname{tr} B^{+,\downarrow}_{r_1,p_2}(v) + \operatorname{tr} B^{-,\downarrow}_{p_1-r_1,p_2}(v)
\end{aligned}$$

and the linear functions on $Q(p_1, r_2)$ by

$$\begin{aligned}
&\tau^{+}_{p_1,r_2}(v) = \operatorname{tr} B^{+,\uparrow}_{p_1,r_2}(v),\ \tau^{-}_{p_1,r_2}(v) = \operatorname{tr} B^{-,\uparrow}_{p_1,p_2-r_2}(v),\\
&\chi_{p_1,r_2}(v) = \operatorname{tr} B^{+,\downarrow}_{p_1,r_2}(v) + \operatorname{tr} B^{-,\downarrow}_{p_1,p_2-r_2}(v)
\end{aligned}$$

for $1 \le r_1 \le p_1 - 1$ and $1 \le r_2 \le p_2 - 1$. These linear functions are symmetric (see [1]).

We also define the linear functions on $Q(r_1, r_2)$ for $(r_1, r_2) \in I$:

$$\begin{aligned}
&\tau^{\uparrow}_{r_1,r_2}(v) = \operatorname{tr} T^{+,\uparrow}_{r_1,r_2}(v),\ \tau^{\downarrow}_{r_1,r_2}(v) = \operatorname{tr} T^{+,\uparrow}_{p_1-r_1,p_2-r_2}(v),\\
&\tau^{\rightarrow}_{r_1,r_2}(v) = \operatorname{tr} T^{-,\uparrow}_{p_1-r_1,r_2}(v),\ \tau^{\leftarrow}_{r_1,r_2}(v) = \operatorname{tr} T^{-,\uparrow}_{r_1,p_2-r_2}(v),\\
&v^{\nearrow}_{r_1,r_2}(v) = \operatorname{tr} T^{+,\downarrow}_{r_1,r_2}(v) + \operatorname{tr} T^{-,\downarrow}_{p_1-r_1,r_2}(v),\ v^{\swarrow}_{r_1,r_2}(v) = \operatorname{tr} T^{-,\downarrow}_{r_1,p_2-r_2}(v) + \operatorname{tr} T^{+,\downarrow}_{p_1-r_1,p_2-r_2}(v),\\
&v^{\nwarrow}_{r_1,r_2}(v) = \operatorname{tr} B^{+,\uparrow}_{r_1,r_2}(v) + \operatorname{tr} B^{-,\uparrow}_{r_1,p_2-r_2}(v),\ v^{\searrow}_{r_1,r_2}(v) = \operatorname{tr} B^{-,\uparrow}_{p_1-r_1,r_2}(v) + \operatorname{tr} B^{+,\uparrow}_{p_1-r_1,p_2-r_2}(v),\\
&\chi_{r_1,r_2}(v) = \operatorname{tr} B^{+,\downarrow}_{r_1,r_2}(v) + \operatorname{tr} B^{-,\downarrow}_{r_1,p_2-r_2}(v) + \operatorname{tr} B^{-,\downarrow}_{p_1-r_1,r_2}(v) + \operatorname{tr} B^{+,\downarrow}_{p_1-r_1,p_2-r_2}(v)
\end{aligned}$$

for $v \in Q(r_1, r_2)$.

Theorem 7.2. (1) *Each of the linear function τ_{p_1,p_2} and τ_{0,p_2} is a basis of the space of symmetric linear functions on $Q(p_1,p_2)$ and $Q(0,p_2)$, respectively.*

(2) *The linear functions*

$$\tau^{\pm}_{r_1,p_2},\ \chi_{r_1,p_2}$$

form a basis of the space of symmetric linear functions on $Q(r_1,p_2)$.

(3) *The linear functions*

$$\tau^{\pm}_{p_1,r_2},\ \chi_{p_1,r_2}$$

form a basis of the space of symmetric linear functions on $Q(p_1,r_2)$.

(4) *The linear functions*

$$\tau^{\bullet}_{r_1,r_2},\ \upsilon^{\circ}_{r_1,r_2},\ \chi_{r_1,r_2},\ \bullet\in\{\uparrow,\downarrow,\leftarrow,\rightarrow\},\ \circ\in\{\nearrow,\nwarrow,\searrow,\swarrow\}$$

form a basis of the space of symmetric linear functions on $Q(r_1,r_2)$ for $i\in I$.

Proof. We only prove (4). By Theorem 6.4, $\tau^{\bullet}_{r_1,r_2}$ with $\bullet\in\{\uparrow,\downarrow,\leftarrow,\rightarrow\}$ is symmetric. For $v,w\in Q(r_1,r_2)$, we can see

$$\begin{aligned}
T^{+,\downarrow}_{r_1,r_2}(vw) &= T^{+,\downarrow}_{r_1,r_2}(v)T^{+,\uparrow}_{r_1,r_2}(w) + T^{-,\rightarrow}_{p_1-r_1,r_2}(v)T^{+,\leftarrow}_{r_1,r_2}(w)\\
&\quad + T^{-,\leftarrow}_{p_1-r_1,r_2}(v)T^{+,\rightarrow}_{r_1,r_2}(w) + T^{+,\uparrow}_{r_1,r_2}(v)T^{+,\uparrow}_{r_1,r_2}(w),\\
T^{-,\downarrow}_{p_1-r_1,r_2}(vw) &= T^{-,\downarrow}_{p_1-r_1,r_2}(v)T^{-,\uparrow}_{p_1-r_1}(w) + T^{+,\rightarrow}_{r_1,r_2}(v)T^{-,\leftarrow}_{p_1-r_1,r_2}(w)\\
&\quad + T^{+,\leftarrow}_{r_1,r_2}(v)T^{-,\rightarrow}_{p_1-r_1,r_2}(w) + T^{-,\uparrow}_{p_1-r_1,r_2}(v)T^{-,\downarrow}_{p_1-r_1,r_2}(w).
\end{aligned}$$

Thus $\upsilon^{\nearrow}_{r_1,r_2}$ is symmetric. Similarly we can see that $\upsilon^{\swarrow}_{r_1,r_2}$ is symmetric. We can also see that

$$\begin{aligned}
B^{+,\uparrow}_{r_1,r_2}(vw) &= B^{+,\uparrow}_{r_1,r_2}(v)T^{+,\uparrow}_{r_1,r_2}(w) + R^{-,\uparrow}_{r_1,p_2-r_2}(v)L^{+,\uparrow}_{r_1,r_2}(w)\\
&\quad + L^{-,\uparrow}_{r_1,p_2-r_2}(v)R^{+,\uparrow}_{r_1,r_2}(w) + T^{+,\uparrow}_{r_1,r_2}(v)B^{+,\uparrow}_{r_1,r_2}(w),\\
B^{-,\uparrow}_{r_1,p_2-r_2}(vw) &= B^{-,\uparrow}_{r_1,p_2-r_2}(v)T^{-,\uparrow}_{r_1,p_2-r_2}(w) + R^{+,\uparrow}_{r_1,r_2}(v)L^{-,\uparrow}_{r_1,p_2-r_2}(w)\\
&\quad + L^{+,\uparrow}_{r_1,r_2}(v)R^{-,\uparrow}_{r_1,p_2-r_2}(w) + T^{-,\uparrow}_{r_1,p_2-r_2}(v)B^{-,\uparrow}_{r_1,p_2-r_2}(w),
\end{aligned}$$

which shows $\upsilon^{\nwarrow}_{r_1,r_2}$ is symmetric. Similarly we can see that $\upsilon^{\searrow}_{r_1,r_2}$ is symmetric.

Now we have

$$\begin{aligned}
B^{+,\downarrow}_{r_1,r_2}(vw) &= B^{+,\downarrow}_{r_1,r_2}(v)T^{+,\uparrow}_{r_1,r_2}(w) + B^{-,\rightarrow}_{p_1-r_1,r_2}(v)T^{+,\leftarrow}_{r_1,r_2}(w)\\
&\quad + B^{-,\leftarrow}_{p_1-r_1,r_2}(v)T^{+,\rightarrow}_{r_1,r_2}(w) + B^{+,\uparrow}_{r_1,r_2}(v)T^{+,\downarrow}_{r_1,r_2}(w)\\
&\quad + R^{-,\downarrow}_{r_1,p_2-r_2}(v)L^{+,\uparrow}_{r_1,r_2}(w) + R^{+,\rightarrow}_{p_1-r_1,p_2-r_2}(v)L^{+,\leftarrow}_{r_1,r_2}(w)\\
&\quad + R^{+,\leftarrow}_{p_1-r_1,p_2-r_2}(v)L^{+,\rightarrow}_{r_1,r_2}(w) + R^{-,\uparrow}_{r_1,p_2-r_2}(v)L^{+,\uparrow}_{r_1,r_2}(w)\\
&\quad + L^{-,\downarrow}_{r_1,p_2-r_2}(v)R^{+,\uparrow}_{r_1,r_2}(w) + L^{+,\rightarrow}_{p_1-r_1,p_2-r_2}(v)R^{+,\leftarrow}_{r_1,r_2}(w)\\
&\quad + L^{+,\leftarrow}_{p_1-r_1,p_2-r_2}(v)R^{+,\rightarrow}_{r_1,r_2}(w) + L^{-,\uparrow}_{r_1,p_2-r_2}(v)R^{+,\downarrow}_{r_1,r_2}(w)\\
&\quad + T^{+,\downarrow}_{r_1,r_2}(v)B^{+,\uparrow}_{r_1,r_2}(w) + T^{-,\rightarrow}_{p_1-r_1,r_2}(v)B^{+,\leftarrow}_{r_1,r_2}(w)\\
&\quad + T^{-,\leftarrow}_{p_1-r_1,r_2}(v)B^{+,\rightarrow}_{r_1,r_2}(w) + T^{+,\uparrow}_{r_1,r_2}(v)B^{+,\downarrow}_{r_1,r_2}(w).
\end{aligned}$$

Substituting $(+,r_1,r_2)$ with $(-,p_1-r_1,r_2)$, $(-,r_1,p_2-r_2)$ and $(+,p_1-r_1,p_2-r_2)$, we can describe $B^{-,\downarrow}_{p_1-r_1,r_2}(vw)$, $B^{-,\downarrow}_{r_1,p_2-r_2}(vw)$ and $B^{+,\downarrow}_{p_1-r_1,p_2-r_2}(vw)$ as linear combinations of traces matrix blocks in the matrix realization of $Q(r_1, r_2)$ for $(r_1, r_2) \in I$. Since the trace is symmetric, we can see that χ_{r_1,r_2} is symmetric. $\square$

7.3. **Integrals and symmetric linear functions.** Recall that the left and right integrals on $\mathfrak{g}_{p_1,p_2}$ is given by

$$\lambda(e_1^{m_1} e_2^{m_2} f_1^{n_1} f_2^{n_2} K^\ell) = \delta_{m_1,p_1-1}\delta_{m_2,p_2-1}\delta_{n_1,p_1-1}\delta_{n_2,p_2-1}\delta_{\ell,p_1-p_2},$$
$$\mu(e_1^{m_1} e_2^{m_2} f_1^{n_1} f_2^{n_2} K^\ell) = \delta_{m_1,p_1-1}\delta_{m_2,p_2-1}\delta_{n_1,p_1-1}\delta_{n_2,p_2-1}\delta_{\ell,p_1+p_2},$$

for $0 \le m_i \le p_i - 1$ and $0 \le \ell \le 2p_1p_2 - 1$. By Proposition 2.2 and Proposition 3.3, we can see

$$\begin{aligned} g^{-1} \rightharpoonup \lambda(e_1^{m_1} e_2^{m_2} f_1^{n_1} f_2^{n_2} K^\ell) &= \mu \leftharpoonup g(e_1^{m_1} e_2^{m_2} f_1^{n_1} f_2^{n_2} K^\ell) \\ &= \delta_{m_1,p_1-1}\delta_{m_2,p_2-1}\delta_{n_1,p_1-1}\delta_{n_2,p_2-1}\delta_{\ell,0}. \end{aligned}$$

The following is the generalization of Lemma 4.1 (see [8]).

Lemma 7.3. *For $1 \le m_i, n_i \le p_i - 1$ with $i = 1, 2$,*

$$[e_i^{m_i}, f_i^{n_i}] = \sum_{j=1}^{\min(m_i,n_i)} e_i^{m_i-j} f_i^{n_i-j} F_j^{m_i,n_i}(K),$$

where $F_j^{m_i,n_i}(z) \in \mathbb{C}[z, z^{-1}]$.

By Lemma 7.3, the term $e_1^{p_1-1} e_2^{p_2-1} f_1^{p_1-1} f_2^{p_2-1}$ appears in $B^{\pm,\downarrow}_{s_1-1,s_2-1}(p_1, p_2, s_1, s_2)$ only for $Q(p_1, p_2)$ and $Q(0, p_2)$. Thus we can see the following.

Proposition 7.4.

$$g^{-1}e(p_1, p_2) \rightharpoonup \lambda = \frac{1}{\Phi^+(r_1, r_2)}\tau_{p_1,p_2},$$
$$g^{-1}e(0, p_2) \rightharpoonup \lambda = \frac{1}{\Phi^-(p_1, p_2)}\tau_{0,p_2}.$$

For $Q(r_1, p_2)$, the term $e_1^{p_1-1} e_2^{p_2-1} f_1^{p_1-1} f_2^{p_2-1}$ appears in

$$B^{+,\uparrow\downarrow}_{s_1-1,s_2-1}(r_1, p_2, s_1, s_2) \text{ and } B^{-,\uparrow\downarrow}_{t_1-1,s_2-1}(p_1, r_2, s_1, s_2)$$

by (4.2.2), (4.2.17) and Lemma 7.3.

Proposition 7.5. (1) *For $1 \le r_1 \le p_1 - 1$, we have*

$$g^{-1}e(r_1, p_2) \rightharpoonup \lambda = \frac{1}{\Phi^+(r_1, p_2)}\left\{\chi_{r_1,p_2} - \Psi_1^+(r_1, p_2)\left(\tau^+_{r_1,p_2} + \tau^-_{r_1,p_2}\right)\right\},$$
$$g^{-1}w(r_1, p_2)^\pm \rightharpoonup \lambda = \frac{1}{\Phi^+(r_1, p_2)}\tau^\pm_{r_1,p_2}.$$

(2) *For $1 \le r_2 \le p_2 - 1$, we have*

$$g^{-1}e(p_1, r_2) \rightharpoonup \lambda = \frac{1}{\Phi^+(p_1, r_2)}\left\{\chi_{p_1,r_2} - \Psi_1^+(p_1, r_2)\left(\tau^+_{p_1,r_2} + \tau^-_{p_1,r_2}\right)\right\},$$
$$g^{-1}w(p_1, r_2)^\pm \rightharpoonup \lambda = \frac{1}{\Phi^+(p_1, r_2)}\tau^\pm_{p_1,r_2}.$$

For $Q(r_1,r_2)$ with $(r_1,r_2)\in I$, we can see that the term $e_1^{p_1-1}e_2^{p_2-1}f_1^{p_1-1}f_2^{p_2-1}$ appears in

$$\begin{aligned}
&T^{+,\uparrow\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2), B^{+,\uparrow\downarrow}_{s_1-1,s_2-1}(r_1,r_2,s_1,s_2),\\
&T^{-,\uparrow\downarrow}_{t_1-1,s_2-1}(p_1-r_1,r_2,t_1,s_2), B^{-,\uparrow\downarrow}_{t_1-1,s_2-1}(p_1-r_1,r_2,t_1,s_2),\\
&T^{-,\uparrow\downarrow}_{s_1-1,t_2-1}(r_1,p_2-r_2,s_1,t_2), B^{+,\uparrow\downarrow}_{s_1-1,s_2-1}(r_1,p_2-r_2,s_1,t_2),\\
&T^{+,\uparrow\downarrow}_{t_1-1,t_2-1}(p_1-r_1,p_2-r_2,t_1,t_2), B^{+,\uparrow\downarrow}_{t_1-1,t_2-1}(p_1-r_1,p_2-r_2,t_1,t_2).
\end{aligned}$$

by (4.3.2), (4.3.8), (4.3.10) and Lemma 7.3. Thus we have the following.

Proposition 7.6. *For $(r_1,r_2)\in I$, we have*

$$\begin{aligned}
&g^{-1}e(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\Big\{\Psi_1^+(r_1,r_2)\Psi_2^+(r_1,r_2)\sum_{\bullet\in\{\uparrow,\downarrow,\leftarrow,\rightarrow\}}\tau^{\bullet}_{r_1,r_2}\\
&-\Psi_2^+(r_1,r_2)\sum_{\bullet\in\{\nearrow,\swarrow\}}\upsilon^{\bullet}_{r_1,r_2}-\Psi_1^+(r_1,r_2)\sum_{\bullet\in\{\nwarrow,\searrow\}}\upsilon^{\bullet}_{r_1,r_2}+\chi_{r_1,r_2}\Big\},\\
&g^{-1}v^{\nearrow}(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\left\{\upsilon^{\nearrow}_{r_1,r_2}-\Psi_1^+(r_1,r_2)\left(\tau^{\uparrow}_{r_1,r_2}+\tau^{\rightarrow}_{r_1,r_2}\right)\right\},\\
&g^{-1}v^{\swarrow}(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\left\{\upsilon^{\swarrow}_{r_1,r_2}-\Psi_1^+(r_1,r_2)\left(\tau^{\downarrow}_{r_1,r_2}+\tau^{\leftarrow}_{r_1,r_2}\right)\right\},\\
&g^{-1}v^{\nwarrow}(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\left\{\upsilon^{\nwarrow}_{r_1,r_2}-\Psi_2^+(r_1,r_2)\left(\tau^{\uparrow}_{r_1,r_2}+\tau^{\leftarrow}_{r_1,r_2}\right)\right\},\\
&g^{-1}v^{\searrow}(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\left\{\upsilon^{\searrow}_{r_1,r_2}-\Psi_2^+(r_1,r_2)\left(\tau^{\downarrow}_{r_1,r_2}+\tau^{\rightarrow}_{r_1,r_2}\right)\right\},\\
&g^{-1}w^{\bullet}(r_1,r_2)\rightharpoonup\lambda=\frac{1}{\Phi^+(r_1,r_2)}\tau^{\bullet}_{r_1,r_2},\ \bullet\in\{\uparrow,\downarrow,\leftarrow,\rightarrow\}.
\end{aligned}$$

Appendix A. q-characters and symmetric linear functions

Let A be a finite-dimensional Hopf algebra. Then the space of q-characters $\mathrm{Ch}(A)$ of A is defined by

$$\begin{aligned}
\mathrm{Ch}(A)&=\{\beta\in A^*\mid \mathrm{Ad}_x(\beta)=\varepsilon(x)\beta\text{ for all }x\in A\}\\
&=\{\beta\in A^*\mid \beta(xy)=\beta(S^2(y)x)\text{ for all }x,y\in A\}
\end{aligned}$$

where $\mathrm{Ad}_a(\beta)=\beta(\sum S(a_1)?a_2)$.

Let $A=\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}}$. Then $g=K^{p_1-p_2}$ is a group-like element and for any $x\in\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}}$ we have $S^2(x)=gxg^{-1}$. It is shown in [5] that the linear function

$$\gamma^{\pm}(r_1,r_2):{\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}}}^*\to\mathbb{C},\ x\mapsto\mathrm{tr}_{X^{\pm}_{r_1,r_2}}(g^{-1}x).$$

for $1\le r_1\le p_1$ and $1\le r_2\le p_2$ belongs to $\mathrm{Ch}(\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}})$.

Note that the map $\theta:\mathrm{Ch}(\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}})\to\mathrm{SLF}(\boldsymbol{\mathfrak{g}}_{\boldsymbol{p_1,p_2}})$ defined by $\beta\mapsto\beta\leftharpoonup g$ is an isomorphism.

Let $\mathbb{P}_{r_1,r_2}$ be the direct sum of four indecomposable projective modules:

$$\mathbb{P}_{r_1,r_2} = \begin{cases} P^+_{r_1,p_2} \oplus P^-_{p_1-r_1,p_2}, & 1 \le r_1 \le p_1 - 1,\ r_2 = p_2, \\ P^+_{p_1,r_2} \oplus P^-_{p_1,p_2-r_2}, & 1 \le r_2 \le p_2 - 1,\ r_1 = p_1, \\ P^+_{r_1,r_2} \oplus P^-_{p_1-r_1,r_2} \oplus P^-_{r_1,p_2-r_2} \oplus P^+_{p_1-r_1,p_2-r_2}, & (r_1,r_2) \in I. \end{cases} \tag{A.0.1}$$

A.1. **q-characters and symmetric linear functions.** Define the linear maps $\sigma_{p_1,r_2} : \mathbb{P}_{p_1,r_2} \to \mathbb{P}_{p_1,r_2}$ and $\sigma_{r_1,p_2} : \mathbb{P}_{r_1,p_2} \to \mathbb{P}_{r_1,p_2}$: the maps σ_{p_1,r_2} and σ_{r_1,p_2} act as zero except

$$\begin{aligned} \sigma_{p_1} &: \mathsf{b}^{\downarrow,\bullet}_{n_1,n_2} \mapsto \alpha^{\uparrow,\bullet}\mathsf{b}^{\uparrow,\bullet}_{n_1,n_2} + \alpha^{\downarrow,\bullet}\mathsf{b}^{\downarrow,\bullet}_{n_1,n_2}, \\ \sigma_{p_2} &: \mathsf{b}^{\downarrow,\circ}_{n_1,n_2} \mapsto \beta^{\uparrow,\circ}\mathsf{b}^{\uparrow,\circ}_{n_1,n_2} + \beta^{\downarrow,\circ}\mathsf{b}^{\downarrow,\circ}_{n_1,n_2}. \end{aligned}$$

where $\bullet \in \{\uparrow, \rightarrow\}$ and $\circ \in \{\downarrow, \leftarrow\}$. In the above equations, $\mathsf{b}^{\downarrow,\uparrow}_{n_1,n_2}$ corresponds to the basis of $P^+_{p_1,r_1}$, $\mathsf{b}^{\downarrow,\rightarrow}_{n_1,n_2}$ corresponds to the basis of $P^-_{p_1,p_2-r_2}$.

We also define the linear map σ_{r_1,r_2} with $(r_1,r_2) \in I$: the map σ_{r_1,r_2} acts as zero except

$$\mathsf{b}^{\downarrow,\bullet}_{n_1,n_2} \mapsto \alpha^{\uparrow,\bullet}\mathsf{b}^{\uparrow,\bullet}_{n_1,n_2} + \alpha^{\downarrow,\bullet}\mathsf{b}^{\downarrow,\bullet}_{n_1,n_2} + \beta^{\uparrow,\bullet}\mathsf{t}^{\uparrow,\bullet}_{n_1,n_2} + \beta^{\downarrow,\bullet}\mathsf{t}^{\downarrow,\bullet}_{n_1,n_2}$$

where $\bullet \in \{\uparrow, \downarrow, \leftarrow, \rightarrow\}$. In the above equation, $\mathsf{b}^{\downarrow,\bullet}_{n_1,n_2}$ for $\bullet =\uparrow$ corresponds to the basis of $P^+_{r_1,r_2}$, for $\bullet =\leftarrow$ corresponds to the basis of $P^-_{r_1,p_2-r_2}$, for $\bullet =\rightarrow$ corresponds to the basis of $P^-_{p_1-r_1,r_2}$ and for $\bullet =\downarrow$ corresponds to the basis of $P^-_{p_1-r_1,p_2-r_2}$.

Then we define $\gamma(r_1,r_2) : x \mapsto \mathrm{tr}_{\mathbb{P}_{r_1,r_2}}(g^{-1}x\sigma_{r_1,r_2})$. In [5], the following is shown.

Proposition A.1 ([5], Proposition 2.3)**.** *The map $\gamma(r_1,r_2)$ belongs to* $\mathrm{Ch}(\mathfrak{g}_{p_1,p_2})$ *if and only if*

$$\begin{aligned} &\alpha^{\uparrow,\uparrow} = \alpha^{\uparrow,\rightarrow},\ \alpha^{\uparrow,\downarrow} = \alpha^{\uparrow,\leftarrow}, \\ &\beta^{\downarrow,\uparrow} = \beta^{\downarrow,\leftarrow},\ \beta^{\downarrow,\downarrow} = \beta^{\downarrow,\rightarrow}, \\ &\beta^{\uparrow,\uparrow} = \beta^{\uparrow,\leftarrow} = \beta^{\uparrow,\rightarrow} = \beta^{\uparrow,\downarrow}. \end{aligned}$$

Theorem A.2 ([5])**.** *The following form a basis of* $\mathrm{Ch}(\mathfrak{g}_{p_1,p_2})$:

$$\begin{aligned} &\gamma^{\pm}(r_1,r_2),\ 1 \le r_1 \le p_1,\ 1 \le r_2 \le p_2, \\ &\gamma^{\nearrow}(r_1,r_2),\ \alpha^{\uparrow,\downarrow} = \alpha^{\downarrow,\bullet} = \beta^{\downarrow,\bullet} = \beta^{\uparrow,\bullet} = 0,\ (r_1,r_2) \in I \cup \{(r_1,p_2)\ |1 \le r_1 \le p_1 - 1\}, \\ &\gamma^{\swarrow}(r_1,r_2),\ \alpha^{\uparrow,\uparrow} = \alpha^{\downarrow,\bullet} = \beta^{\downarrow,\bullet} = \beta^{\uparrow,\bullet} = 0,\ (r_1,r_2) \in I, \\ &\gamma^{\nwarrow}(r_1,r_2),\ \alpha^{\downarrow,\bullet} = \alpha^{\uparrow,\bullet} = \beta^{\downarrow,\downarrow} = \beta^{\uparrow,\bullet} = 0,\ (r_1,r_2) \in I, \\ &\gamma^{\searrow}(r_1,r_2), \alpha^{\downarrow,\bullet} = \alpha^{\uparrow,\bullet} = \beta^{\downarrow,\uparrow} = \beta^{\uparrow,\bullet} = 0,\ (r_1,r_2) \in I \cup \{(p_1,r_2)\ |1 \le r_2 \le p_2 - 1\}, \\ &\gamma^{\downarrow,\downarrow}(r_1,r_2),\ \alpha^{\downarrow,\bullet} = \alpha^{\uparrow,\bullet} = \beta^{\downarrow,\bullet} = 0,\ (r_1,r_2) \in I. \end{aligned}$$

Then by this isomorphism and Table 1, we have

Theorem A.3. *We have*

$$\theta(\gamma^{+}(p_1,p_2)) = \tau_{p_1,p_2},\ \theta(\gamma^{-}(p_1,p_2)) = \tau_{0,r_2},$$

$$\theta(\gamma^{+}(r_1,p_2)) = \tau^{+}_{r_1,p_2},\ \theta(\gamma^{-}(p_1-r_1,p_2) = \tau^{-}_{r_1,p_2},\ \theta(\gamma^{\nearrow}(r_1,p_2)) = \chi_{r_1,p_2}, 1 \leq r_1 \leq p_1-1,$$

$$\theta(\gamma^{+}(p_1,r_2)) = \tau^{+}(p_1,r_2),\ \theta(\gamma^{-}(p_1,p_2-r_2)) = \tau^{-}(p_1,r_2),$$

$$\theta(\gamma^{\searrow}(p_1,r_2)) = \chi_{p_1,r_2},\ 1 \leq r_2 \leq p_2-1,$$

$$\theta(\gamma^{+}(r_1,r_2)) = \tau^{\uparrow}_{r_1,r_2},\ \theta(\gamma^{-}(r_1,p_2-r_2)) = \tau^{\leftarrow}_{r_1,p_2-r_2},$$

$$\theta(\gamma^{-}(p_1-r_1,r_2)) = \tau^{\rightarrow}_{p_1-r_1,r_2},\ \theta(\gamma^{+}(p_1-r_1,p_2-r_2)) = \tau^{\downarrow}_{p_1-r_1,p_2-r_2},$$

$$\theta(\gamma^{\nearrow}(r_1,r_2)) = \upsilon^{\nearrow}_{r_1,r_2},\ \theta(\gamma^{\swarrow}(r_1,r_2)) = \upsilon^{\swarrow}_{r_1,r_2},$$

$$\theta(\gamma^{\nwarrow}(r_1,r_2)) = \upsilon^{\nwarrow}_{r_1,r_2},\ \theta(\gamma^{\searrow}(r_1,r_2)) = \upsilon^{\searrow}_{r_1,r_2},$$

$$\theta(\gamma^{\downarrow,\downarrow}(r_1,r_2)) = \chi_{r_1,r_2},\ (r_1,r_2) \in I.$$

REFERENCES

[1] Y. Arike, A construction of symmetric linear functions on $\overline{U}_q(sl_2)$, preprint, to appear in Osaka. J. Math.

[2] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Yu. Tipunin, Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center, *Commun. Math. Phys.* **265** (2006), 47-93.

[3] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Yu. Tipunin, Kazhdan-Lustig correspondence for the representation category of the triplet W-algebra in logarithmic CFT, *Theor. Math. Phys.* **148** (2006), 1210-1235.

[4] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Yu. Tipunin, Logarithmic extensions of minimal models: characters and modular transformations, *Nucl. Phys. B.* **757** (2006), 303-343.

[5] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov and I. Yu. Tipunin, Kazhdan-Lusztig-dual quantum group for logarithmic extensions of Virasoso minimal models, *J. Math. Phys.* **48**, 032303, (2007),

[6] C. Kassel, *Quantum Groups*, Graduate Text in Mathematics. **155**, Springer-Verlag, New York (1995).

[7] H. Kondo and Y. Saito, Indecomposable decomposition of tensor products of modules over the restricted quantum universal enveloping algebra associated to $\mathfrak{sl}_2$, preprint, arXiv:0901.4221v2.

[8] G. Lusztig, Quantum deformations of certain simple modules over enveloping algebras, *Adv. in Math.* **70**, (1988), no. 2, 237-249.

[9] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Conf. Series in Math. 82, Amer. Math. Soc., Providence, RI, (1993).

[10] K. Nagatomo and A. Tsuchiya, The triplet vertex operator algebra $W(p)$ and the restricted quantum group at root of unity, preprint, arXiv:0902.4607.

[11] D. E. Radford, The trace function and Hopf algebras, *J. Algebra.* **163** (1994), 583-622.

[12] R. Suter, Modules over $\mathfrak{U}_q(\mathfrak{sl}_2)$, *Commun. Math. Phys.* **163** (1994), 359-393.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: `y-arike@cr.math.sci.osaka-u.ac.jp`